## Predicting the chemical shift of proton/s on a sp<sup>3</sup> carbon in a <sup>1</sup>H NMR spectrum

Step one: Find the all chemically non-equivalent proton/s.

When predicting chemical shifts, you consider the chemically non-equivalent proton/s one at a time.

Step two: Find the starting point for each chemically non-equivalent proton/s.

The starting point is dependent upon if the proton/s are attached to a methyl, a methene or a methine.

If the protons are attached to a methyl (XCH<sub>3</sub>) the starting point is 0.9 ppm. If the protons are attached to a methene (XCH<sub>2</sub>X) the starting point is 1.3 ppm. If the proton is attached to a methine ( $X_3$ CH) the starting point is 1.5 ppm.

X can represent anything but a H or D.

Step three: What is X?

Identify all groups (X) that are directly bonded to the carbon atom that bears the proton/s of interest. Be carful with esters because they have two sides, the carbonyl side and the acyoxy side.

Group (X)	Increment (ppm )
—CH₃ (or any alkyl group)	0.0
—CH=CH₂ (or any alkene group)	0.9
—C≡CR, —C≡CH	0.9
—SR, —SH	1.2
O O II II — C—OH	1,.2


Group (X)	Increment (ppm )
—C≡N	1.2
О О П П П П П П П П П П П П П П П П П П	1.2
—NH <sub>2</sub> , —NHR, —NR <sub>2</sub>	1.2
	1.3
—Ph	1.4
L	

Group (X)	Increment (ppm )
—Br	1.8
—CI	2.1
OR,OH	2.4
O II —O—C—R	2.8
—OPh	3.0
-	3.2

Step four: Add the values together.

When predicting chemical shifts, you need to watch for *remote* (not geminal) electronegative group/s. If a substituent has a remote electronegative group, an asterisk (\*) is add to the chemical shift increment indicating that the chemical shift increment will be increased. Adding the starting point and all the X values together (including the asterisk) will provide the approximate chemical shift of the proton of interest.

## What is the predicted chemical shift ( $\delta$ ) of proton Ha?



## What is the predicted chemical shift ( $\delta$ ) of proton Hb?

$$FH_{2}CH_{2}C-\overset{\downarrow}{C}-O-\overset{\downarrow}{C}-O-\overset{\downarrow}{C}-CH_{3}$$

$$XCH_{2}X \text{ starting point} \qquad \qquad \delta \text{ starting point} \qquad \qquad 1.3 \text{ ppm}$$

$$x \text{ is a alkyl group with a remote electronegative atom} \qquad \qquad FH_{2}CH_{2}C -\overset{\downarrow}{C}-\overset{\downarrow}{$$

Predicted
Chemical Shift

4.1\* ppm

Thank-you Dr. J. R Walker at Oregon State University for the foundation of this document