

HW1

Math 344

Week One

1 Computations

1. Let $A = \{a, b, c\}$.

- (a) Write down all functions from A to itself, until you have convinced yourself that you *could* write down all such functions. How many are there? (So the answer to this question is a number.) Here is an example of a function from A to itself:

$$\begin{aligned}a &\mapsto b \\ b &\mapsto c \\ c &\mapsto a.\end{aligned}$$

- (b) Write down all functions f from A to itself with the property that $f \circ f$ is the identity function on A .

- (c) Write down all functions from A to $\{1\}$.

2. Let $B = \{0, 1, 2, 3\}$. Let C be the following set:

$$C = \{(a, b) \in B \times B \mid \text{both } a \text{ and } b \text{ are odd}\}.$$

Write down all elements of C .

2 Proofs

Here are two example problems, with solutions.

Example. For any integers a, b , define $a \otimes b = |ab|$. Prove $\otimes$ is an operation on $\mathbb{Z}$.

Proof. Choose $a, b \in \mathbb{Z}$. Note that if a and b have the same sign, then $a \otimes b = ab$, which is an integer, and if a and b have different signs, then $a \otimes b = -ab$, which is also an integer. Thus, we see that $\otimes$ satisfies the definition of operation. $\square$

Here are some comments about the example.

- Literally everything in the solution is part of a complete sentence.
- Before I manipulate a, b at you, I tell you specifically what they are. (This is required, or I will think to myself “who are these people a and b ? I don’t remember being introduced to them”).
- I don’t restate the definition of operation, because that is part of the set of common knowledge for this class (ie, you are writing these proofs *for me*, and I know all the definitions.)

The “easier” version of this type of question is when you must show a statement is false—you need only provide a counterexample. Often you can even skip introductions, because you will not need variables!

Example. For any $a, b \in \mathbb{Z}_{\neq 0}$, define $a \bullet b = a/b$. Prove that $\bullet$ is not an operation on $\mathbb{Z}_{\neq 0}$.

Proof. Note that 1, 2 are nonzero integers but $1 \bullet 2 = 1/2$, which is not an integer. Therefore, we see that $\bullet$ is not an operation on $\mathbb{Z}_{\neq 0}$. $\square$

Here are some exercises. Fill in the blanks *with complete sentences*.

1. For any integers a, b , define $a \otimes b = a - b$. Prove $\otimes$ is an operation on $\mathbb{Z}$.

Proof. Choose $a, b \in \mathbb{Z}$. Note that $a \otimes b = a - b$, which is also an integer. $\square$

2. For any positive integers a, b , define $a \bullet b = a - b$. Prove $\bullet$ is not an operation on $\mathbb{Z}_{>0}$.

Proof. $\square$. Therefore, we see that $\bullet$ is not an operation on $\mathbb{Z}_{>0}$. $\square$

3. For any $a, b \in \mathbb{Q}_{\neq 0}$, define $a \odot b = a/b$. Then $\odot$ is an operation on $\mathbb{Q}_{\neq 0}$.

Proof. Choose any $a, b \in \mathbb{Q}_{\neq 0}$. $\square$. We see that

$$a \odot b = \frac{c}{d} \odot \frac{e}{f} = \frac{\frac{c}{d}}{\frac{e}{f}} = \frac{cf}{ed}.$$

Since c, d, e, f are nonzero as we noted above, we see that ce, df are nonzero, so that $\frac{cf}{ed} \in \mathbb{Q}_{\neq 0}$. Therefore, we conclude that $\odot$ is an operation on $\mathbb{Q}_{\neq 0}$. $\square$

4. Consider for a moment the operation of multiplication on the integers. Prove that this operation admits an identity.

Proof. $\square$. Thus, we see 1 is the identity element for multiplication on the integers. $\square$

Finally, here are opportunities to write complete proofs:

1. In [Exercise \(2\)](#), we defined the set C . For any $(a, b), (c, d) \in C$, define $(a, b) \oplus (c, d) = (a + c, b + d)$. Prove that $\oplus$ is not an operation on C .
2. Suppose that S, T are sets. Prove that $S \cap T = S$ if and only if $S \subseteq T$.
3. Suppose that m is an integer. Prove:

$$\text{if } 20 \mid m, \text{ then } 4 \mid m.$$