

THE DPG-STAR METHOD

LESZEK DEMKOWICZ, JAY GOPALAKRISHNAN, AND BRENDAN KEITH

ABSTRACT. This article introduces the DPG-star (from now on, denoted DPG*) finite element method. It is a method that is in some sense dual to the discontinuous Petrov–Galerkin (DPG) method. The DPG methodology can be viewed as a means to solve an overdetermined discretization of a boundary value problem. In the same vein, the DPG* methodology is a means to solve an underdetermined discretization. These two viewpoints are developed by embedding the same operator equation into two different saddle-point problems. The analyses of the two problems have many common elements. Comparison to other methods in the literature round out the newly garnered perspective. Notably, DPG* and DPG methods can be seen as generalizations of \mathcal{LL}^* and least-squares methods, respectively. *A priori* error analysis and *a posteriori* error control for the DPG* method are considered in detail. Reports of several numerical experiments are provided which demonstrate the essential features of the new method. A notable difference between the results from the DPG* and DPG analyses is that the convergence rates of the former are limited by the regularity of an extraneous Lagrange multiplier variable.

1. INTRODUCTION

The *ideal Discontinuous Petrov–Galerkin (DPG) Method with Optimal Test Functions* [18, 20] admits three interpretations [22]. First, it can be viewed as a Petrov–Galerkin (PG) discretization in which *optimal* test functions are computed on the fly. Here, the word “optimal” refers to the fact that the test functions realize the supremum in the discrete inf-sup stability condition and, therefore, the PG discretization automatically inherits the stability of the continuous method. The DPG method can also be viewed as a minimum residual method in which the residual is measured in a dual norm implied by an underlying test norm. Finally, the DPG method can be viewed also as a mixed method [16] wherein one simultaneously solves for the Riesz representation of the residual — the so-called *error representation function* — and the approximate solution. All three equivalent interpretations involve the inversion of a Riesz operator on the test space which, in general, cannot be done exactly and has to be approximated. This naturally leads to the introduction of an *enriched* or *search* test space — having dimension larger than that of the latent trial space — and a discretized Riesz operator. In this way, the corresponding *practical DPG method* retains its three interpretations, although now with *approximate* optimal test functions, an *approximate* residual, and an *approximate* error representation function [35, 43, 11].

2010 *Mathematics Subject Classification.* 65J05, 65N12, 65N15, 65N30.

Key words and phrases. DPG method, ultraweak formulation, a priori, a posteriori, duality.

Acknowledgments. This work was partially supported with grants by NSF (DMS-1418822), AFOSR (FA9550-17-1-0090), and ONR (N00014-15-1-2496). Part of this work was done while the second author was at The University of Texas at Austin on a J. Tinsley Oden Faculty Fellowship. The third author was additionally supported in part by the 2017 Graduate School University Graduate Continuing Fellowship at The University of Texas at Austin. The authors express their gratitude to Nathan V. Roberts and Socratis Petrides for their assistance during the numerical experiments with certain features of the Camellia and *hp2D* software packages.

In the DPG method, the word “discontinuous” corresponds to the use of discontinuous, but conforming, test functions (from broken spaces) which make the whole methodology computationally efficient. These broken spaces also naturally lead to (hybridized) interface solution variables. Broken space formulations provide a foundation for the DPG methodology and can be developed for any well-posed variational formulation [11].

Originally motivated by the duality theory in [39], in this article, the DPG methodology is expanded on by reconsidering how a given operator equation can be embedded into a DPG-type saddle-point problem. In turn, the minimum residual principle underpinning DPG methods can be discarded for a minimum norm principle and a dual class of methods (i.e. DPG* methods¹) can be introduced. Ultimately, an entire class of stable DPG-type mixed methods can be proposed, simply by changing loads in the saddle-point problem and the interpretation of the corresponding solution variables. This broad perspective can help relate several different methods, including weakly conforming least squares methods [28] and \mathcal{LL}^* methods [10] to the existing DPG theory. In addition to being a mathematical curiosity, the DPG* framework rounds out the *a priori* error analysis of DPG methods in weak norms considered in previous articles (e.g., [6, 32, 33]), is instrumental in goal-oriented *a posteriori* error analysis of DPG methods [39], and may be used to design new adjoint methods with DPG [38], such as for the solution of optimal control problems.

This article provides a number of important *a priori* error analysis results for DPG* methods. Unlike standard DPG methods, the convergence rate of a DPG* solution is not controlled solely by the regularity of the solution itself, but instead also by the regularity of a Lagrange multiplier variable found in the corresponding saddle-point formulation. This article also expands on the recent *a posteriori* error estimation theory first introduced in [39] and re-establishes much of Repin, Sauter, and Smolianski’s abstract *a posteriori* theory for mixed methods [45] in the present context. In addition, it includes several standard numerical examples to verify the theory for DPG* methods, including one example employing *hp*-adaptive mesh refinement. This work is part of the PhD thesis [38].

2. THE DPG AND DPG* METHODS

2.1. Operator equations. Central to this paper are the twin relatives of the operator equation

$$(1) \quad Bu = \ell,$$

given in (2) and (3) below. Here $B : U \rightarrow V'$ is a bounded linear operator from a Hilbert space U to the dual of a Hilbert space V , $\ell \in V'$ is given, and $u \in U$ is to be found. All spaces here are over \mathbb{R} , the real field. In any Hilbert space X , the action of a functional $E \in X'$ on $x \in X$ is denoted by $\langle E, x \rangle_X$. When the space is clear from context, we also use $E(x)$ to denote the same number. Let $B' : V \rightarrow U'$ be the dual of B defined by $\langle B'v, u \rangle_U = \langle Bu, v \rangle_V$ for all $u \in U$ and $v \in V$. The reason for using $'$ instead of $*$ to denote the dual operator will become evident when a different, but related, notion of duality is introduced in Section 2.3.²

¹DPG* methods are distinct from the saddle point least squares methods [2] which have separately been contenders for being named “dual” to DPG methods.

²Therefore, the asterisk in the DPG* method is evocative of the connections to the \mathcal{LL}^* method [10].

The two reformulations are as follows.

$$\begin{aligned}
 (2) \quad & \text{Find } u \in U \text{ and } \varepsilon \in V \text{ satisfying} & \begin{cases} \mathcal{R}_V \varepsilon + Bu = \ell, \\ B' \varepsilon = 0. \end{cases} \\
 (3) \quad & \text{Find } u \in U \text{ and } \lambda \in V \text{ satisfying} & \begin{cases} \mathcal{R}_U u - B' \lambda = 0, \\ Bu = \ell. \end{cases}
 \end{aligned}$$

Here $\mathcal{R}_V : V \rightarrow V'$ is the Riesz operator acting on V , defined using the inner product $(\cdot, \cdot)_V$ by $(\mathcal{R}_V v)(\nu) = (v, \nu)_V$ for all $v, \nu \in V$. The Riesz operator \mathcal{R}_U is defined similarly. It is immediate that if u solves (1), then with $\varepsilon = 0$ it solves (2), revealing a relationship between (2) and (1). The relationship between (3) and (1) is also easy to guess: any solution (u, λ) of (3) is such that the u component solves (1). We shall see below that, even though related, these formulations are not fully equivalent to (1). The formulation (2) is the one on which the DPG method is based. The formulation (3), when discretized, results in the new DPG* method, as we shall see.

Formulations (2) and (3) are structurally similar, differing mainly in the position where the load ℓ is placed. Due to the structural similarity, both formulations can be viewed at once as instantiations of the following general saddle-point problem

$$(4) \quad \text{Find } v \in \mathcal{V} \text{ and } w \in \mathcal{U} \text{ satisfying} \quad \begin{cases} \mathcal{R}_V v + \mathcal{B}w = F, \\ \mathcal{B}'v = G, \end{cases}$$

on some Hilbert spaces \mathcal{U} and \mathcal{V} , some bounded linear operator $\mathcal{B} : \mathcal{U} \rightarrow \mathcal{V}'$, and some given functionals $F \in \mathcal{V}'$ and $G \in \mathcal{U}'$. Indeed, with

$$\mathcal{V} = V, \mathcal{U} = U, \mathcal{B} = B, F = \ell, G = 0,$$

we obtain (2). If instead, we set

$$\mathcal{V} = U, \mathcal{U} = V, \mathcal{B} = -B', F = 0, G = -\ell,$$

then we obtain (3). Admittedly, the alternative mixed form obtained by exchanging \mathcal{B} and \mathcal{B}' in (4) is more natural for studying the DPG* method and even aligns with the standard notations in mixed method theory [5]. Yet, we have chosen to work with (4) to facilitate comparison with existing DPG literature where the form of (4) is more natural.

We proceed under the assumption that \mathcal{B} is bounded below, i.e., there is a $\gamma > 0$ such that

$$(5) \quad \|\mathcal{B}\mu\|_{\mathcal{V}'} \geq \gamma \|\mu\|_{\mathcal{U}}, \quad \forall \mu \in \mathcal{U}.$$

Note that the maximum of all such γ is simply $\|\mathcal{B}^{-1}\|$. Under this assumption, the mixed system (4) has a unique solution for any $F \in \mathcal{V}'$ and $G \in \mathcal{U}'$ (see e.g. [5]). Obviously (5) can also be written out as an inf-sup condition. Here and throughout, for any Banach space X , the right annihilator of a subset $Y \subseteq X$ and the left annihilator of a $Z \subseteq X'$ are defined by

$$(6) \quad Y^\perp = \{E \in X' : \langle E, y \rangle_X = 0 \text{ for all } y \in Y\},$$

$$(7) \quad {}^\perp Z = \{x \in X : \langle E, x \rangle_X = 0 \text{ for all } E \in Z\}.$$

Recall that if $Y \subseteq X$ is a closed subspace, $\overline{Y} = Y$, then Y^\perp is isomorphic to $(X/Y)'$.

Now consider the mixed system (4) when $G = 0$ and the related problem of finding $w \in \mathcal{U}$ satisfying

$$(8) \quad \mathcal{B}w = F.$$

The regularizing effect of the saddle-point formulation above is already evident: while (4) is always solvable under (5), the related problem (8) is solvable *provided* F satisfies the compatibility condition $F \in (\text{Null } \mathcal{B}')^\perp$. To reiterate the above observation that (8) is not fully equivalent to (4), we may view (8) as an *overdetermined system*. Overdetermined systems are solvable only if they are consistent, i.e., have compatible data. Irrespective of the data, what the mixed system (4) solves can be seen by eliminating v (and recalling that $G = 0$), namely

$$(9) \quad \mathcal{B}' \mathcal{R}_\mathcal{V}^{-1} \mathcal{B} w = \mathcal{B}' \mathcal{R}_\mathcal{V}^{-1} F.$$

Equation (9) can be immediately identified with what is referred to as a “normal equation” in linear algebra. This is a regularized version of (8). Indeed, whenever (8) has a solution, it must be unique due to (5), and that unique solution is recovered by (9). However, (9) has a unique solution even when (8) does not.

Next, considering the case of $F = 0$, we may likewise argue that the mixed system (4) also helps us solve *underdetermined systems*. Under the same assumption (5), consider

$$(10) \quad \mathcal{B}' v = G.$$

Assumption (5) implies that \mathcal{B}' is surjective, so (10) is always solvable, but its solution need not be unique in general. Thus, (10) may be viewed as an example of an underdetermined system. Similar to (9), the solution variable v can be readily eliminated from (4) (now recalling that $F = 0$):

$$(11) \quad \mathcal{B}' \mathcal{R}_\mathcal{V}^{-1} \mathcal{B} w = -G.$$

This equation is in correspondence with a different normal equation (one of the second type [3]). Notice that the left-hand side operator $\mathcal{B}' \mathcal{R}_\mathcal{V}^{-1} \mathcal{B} : \mathcal{U} \rightarrow \mathcal{U}'$ is the same in both (9) and (11) and that the solution v in (11) can be recovered by the relationship $v = -\mathcal{R}_\mathcal{V}^{-1} \mathcal{B} w$.

To reconsider how the mixed system (4) converts (10) into a uniquely solvable problem, we use orthogonal complements in Hilbert spaces, which we distinguish from the annihilators in (6) and (7) by placing the symbol \perp as a subscript. Thus, while $(\text{Null } \mathcal{B}')^\perp$ is a subspace of \mathcal{V}' , the notation $(\text{Null } \mathcal{B}')_\perp$ indicates the subspace of \mathcal{V} defined by

$$(\text{Null } \mathcal{B}')_\perp = \{v \in \mathcal{V} : (v, \nu_0)_\mathcal{V} = 0, \forall \nu_0 \in \text{Null } \mathcal{B}'\}.$$

One may then decompose any solution of (10) into \mathcal{V} -orthogonal components:

$$(12) \quad v = v_0 + v_\perp, \quad v_0 \in \text{Null } \mathcal{B}', \quad v_\perp \in (\text{Null } \mathcal{B}')_\perp.$$

Observe that

$$(13) \quad (\text{Null } \mathcal{B}')^\perp = \mathcal{R}_\mathcal{V}(\text{Null } \mathcal{B}')_\perp.$$

Since $F = 0$, testing the first equation of (4) with v_0 , we find that what (4) selects as its unique solution v is in fact simply v_\perp .

Returning to the case of general F and G , we collect a few identities in the next result. First, note that one may also decompose F into orthogonal components:

$$(14) \quad F = F^0 + F^\perp, \quad F^0 \in \mathcal{R}_\mathcal{V}(\text{Null } \mathcal{B}'), \quad F^\perp \in \mathcal{R}_\mathcal{V}(\text{Null } \mathcal{B}')_\perp = (\text{Null } \mathcal{B}')^\perp.$$

Second, note that when (5) holds, $\|\mu\|_\mathcal{U} = \|\mathcal{B}\mu\|_\mathcal{V}$ generates an equivalent norm on \mathcal{U} , $\|\mathcal{B}^{-1}\|^{-1}\|\mu\|_\mathcal{U} \leq \|\mu\|_\mathcal{U} \leq \|\mathcal{B}\|\|\mu\|_\mathcal{U}$, and we may define

$$(15) \quad \|G\|_{\mathcal{U}'} = \sup_{0 \neq \mu \in \mathcal{U}} \frac{\langle G, \mu \rangle_{\mathcal{U}'}}{\|\mu\|_\mathcal{U}}.$$

Proposition 2.1. *Suppose $F \in \mathcal{V}'$, $G \in \mathcal{U}'$, $v \in \mathcal{V}$ and $w \in \mathcal{U}$ solve (4) and let v_0 and v_\perp be the unique components of the decomposition of v in (12). Similarly, let F^0 and F^\perp be the unique components of the decomposition of F in (14). Then the following identities hold:*

$$(16) \quad \|v_0\|_{\mathcal{V}}^2 + \|\mathcal{R}_{\mathcal{V}}v_\perp + \mathcal{B}w\|_{\mathcal{V}'}^2 = \|F\|_{\mathcal{V}'}^2,$$

$$(17) \quad \|v_0\|_{\mathcal{V}}^2 + \|\mathcal{B}w\|_{\mathcal{V}'}^2 = \|F - \mathcal{R}_{\mathcal{V}}v_\perp\|_{\mathcal{V}'}^2.$$

Moreover, $v_0 = \mathcal{R}_{\mathcal{V}}^{-1}F^0$ and

$$(18) \quad \|v_0\|_{\mathcal{V}} = \|F^0\|_{\mathcal{V}'},$$

$$(19) \quad \|\mathcal{B}w\|_{\mathcal{V}'} = \|F^\perp - \mathcal{R}_{\mathcal{V}}v_\perp\|_{\mathcal{V}'}.$$

If in addition, (5) holds, then for any $F \in \mathcal{V}'$, $G \in \mathcal{U}'$, there is a unique $v \in \mathcal{V}$ and $w \in \mathcal{U}$ satisfying (4) and the following identities hold:

$$(20) \quad \|v_\perp\|_{\mathcal{V}} = \|G\|_{\mathcal{U}},$$

$$(21) \quad \|v\|_{\mathcal{V}}^2 + \|w\|_{\mathcal{U}}^2 = \|F - \mathcal{R}_{\mathcal{V}}v_\perp\|_{\mathcal{V}'}^2 + \|G\|_{\mathcal{U}}^2.$$

If in addition, either $F \in (\text{Null } \mathcal{B}')^\perp$ or \mathcal{B} is a bijection, then $v_0 = 0$ and

$$(22) \quad \|v\|_{\mathcal{V}} = \|G\|_{\mathcal{U}}.$$

Proof. For any $\nu_0 \in \text{Null } \mathcal{B}'$, we have $(\mathcal{R}_{\mathcal{V}}^{-1}\mathcal{B}w, \nu_0)_{\mathcal{V}} = \langle \mathcal{B}w, \nu_0 \rangle_{\mathcal{V}} = \langle \mathcal{B}'\nu_0, w \rangle_{\mathcal{U}} = 0$. Hence $\mathcal{R}_{\mathcal{V}}^{-1}\mathcal{B}w$ is in $(\text{Null } \mathcal{B}')^\perp$. Therefore, when the first equation of (4) is rewritten as

$$(23) \quad v_0 + (v_\perp + \mathcal{R}_{\mathcal{V}}^{-1}\mathcal{B}w) = \mathcal{R}_{\mathcal{V}}^{-1}F,$$

an application of the Pythagorean theorem gives (16). Rewriting (23) as $v_0 + \mathcal{R}_{\mathcal{V}}^{-1}\mathcal{B}w = \mathcal{R}_{\mathcal{V}}^{-1}F - v_\perp$, and applying the Pythagorean theorem again, we obtain (17). Rewriting (23) instead as

$$v_0 - \mathcal{R}_{\mathcal{V}}^{-1}F^0 = \mathcal{R}_{\mathcal{V}}^{-1}F^\perp - (v_\perp + \mathcal{R}_{\mathcal{V}}^{-1}\mathcal{B}w),$$

we note that $v_0 = \mathcal{R}_{\mathcal{V}}^{-1}F^0$ and $\mathcal{R}_{\mathcal{V}}^{-1}F^\perp = v_\perp + \mathcal{R}_{\mathcal{V}}^{-1}\mathcal{B}w$, by orthogonality. Equations (18) and (19) are now obvious.

Next, if (5) holds, then standard mixed theory [5] gives existence of a unique $(v, w) \in \mathcal{V} \times \mathcal{U}$, and $\|\cdot\|_{\mathcal{U}}$ is an equivalent norm on \mathcal{U} . To prove (20), we begin by noting that the isometry induced by $\mathcal{R}_{\mathcal{V}}$ implies

$$\|v_\perp\|_{\mathcal{V}} = \sup_{\nu_\perp \in (\text{Null } \mathcal{B}')^\perp} \frac{(\nu_\perp, v_\perp)_{\mathcal{V}}}{\|\nu_\perp\|_{\mathcal{V}}} = \sup_{\nu_\perp \in (\text{Null } \mathcal{B}')^\perp} \frac{\langle \mathcal{R}_{\mathcal{V}}\nu_\perp, v_\perp \rangle_{\mathcal{V}}}{\|\mathcal{R}_{\mathcal{V}}\nu_\perp\|_{\mathcal{V}'}} = \sup_{E^\perp \in \mathcal{R}_{\mathcal{V}}(\text{Null } \mathcal{B}')^\perp} \frac{\langle E^\perp, v_\perp \rangle_{\mathcal{V}}}{\|E^\perp\|_{\mathcal{V}'}}.$$

Here and throughout, supremums over spaces are only taken over nonzero elements of the space. Again, from the identity $\text{Range } \mathcal{B} = (\text{Null } \mathcal{B}')^\perp$ and (13), we conclude that

$$\|v_\perp\|_{\mathcal{V}} = \sup_{E^\perp \in \text{Range } \mathcal{B}} \frac{\langle E^\perp, v_\perp \rangle_{\mathcal{V}}}{\|E^\perp\|_{\mathcal{V}'}} = \sup_{\mu \in \mathcal{U}} \frac{\langle \mathcal{B}\mu, v_\perp \rangle_{\mathcal{V}}}{\|\mathcal{B}\mu\|_{\mathcal{V}'}} = \sup_{\mu \in \mathcal{U}} \frac{\langle \mu, \mathcal{B}'v_\perp \rangle_{\mathcal{V}}}{\|\mu\|_{\mathcal{U}}}.$$

Thus, (20) follows after using the second equation in (4), namely $G = \mathcal{B}'(v_0 + v_\perp) = \mathcal{B}'v_\perp$. Identity (21) now follows by squaring both sides of (20) and adding it to (17).

Finally, when \mathcal{B} is a bijection or $F \in (\text{Null } \mathcal{B}')^\perp$, we conclude that $F^0 = 0$. Therefore, $v_0 = 0$ and (22) follows from (20). \square

Identities like (21) have often been referred to by the name *hypercircle identities* [45] and their use in *a posteriori* error estimation is now standard. We shall return to this in Section 4.

2.2. Forms and discretization. It is traditional to write mixed systems using a bilinear form defined by

$$(24) \quad b(\mu, \nu) = \langle \mathcal{B}\mu, \nu \rangle_{\mathcal{V}}$$

for all $\mu \in \mathcal{U}, \nu \in \mathcal{V}$. In terms of b , the mixed problem (4) is to find $v \in \mathcal{V}$ and $w \in \mathcal{U}$ satisfying

$$(25) \quad \begin{cases} (v, \nu)_{\mathcal{V}} + b(w, \nu) = F(\nu), & \forall \nu \in \mathcal{V}, \\ b(\mu, v) = G(\mu), & \forall \mu \in \mathcal{U}. \end{cases}$$

Suppose b arises from a weak formulation of a PDE on a domain $\Omega \subseteq \mathbb{R}^d$, which is partitioned into a mesh Ω_h of finitely many open connected elements K with Lipschitz boundaries ∂K , such that $\bar{\Omega}$ is the union of the closures of all mesh elements K in Ω_h . In this scenario, if there are Hilbert spaces $\mathcal{V}(K)$ on each mesh element K such that

$$(26) \quad \mathcal{V} = \prod_{K \in \Omega_h} \mathcal{V}(K),$$

then the system (25), in the case $G = 0$, is called a *DPG formulation*. In the case $F = 0$, it is called a *DPG* formulation*. Spaces of the form (26) are called *broken spaces* [11].

A discrete method based on (25) would require a pair of discrete finite-dimensional spaces $\mathcal{U}_h \subseteq \mathcal{U}$ and $\mathcal{V}_h \subseteq \mathcal{V}$, not necessarily of the same dimension. The discrete problem would then read as the problem of finding $v_h \in \mathcal{V}_h$ and $w_h \in \mathcal{U}_h$ satisfying

$$(27) \quad \begin{cases} (v_h, \nu)_{\mathcal{V}} + b(w_h, \nu) = F(\nu), & \forall \nu \in \mathcal{V}_h, \\ b(\mu, v_h) = G(\mu), & \forall \mu \in \mathcal{U}_h. \end{cases}$$

When \mathcal{V} is a broken space of the form (26), \mathcal{V}_h can be chosen to consist of functions with no continuity constraints across mesh element interfaces. Then the case $G = 0$ delivers *DPG methods* and the case $F = 0$ delivers *DPG* methods*. In both cases, we must typically find \mathcal{V}_h with $\dim(\mathcal{V}_h) > \dim(\mathcal{U}_h)$ with provable discrete stability.

A key feature of (27) is that the top left form, $(v, \nu)_{\mathcal{V}}$, being an inner product, is always coercive. Hence the discrete stability of (27) is guaranteed solely by a discrete inf-sup condition, which is often *easy to obtain* in practice since we can increase $\dim(\mathcal{V}_h)$ without violating the coercivity of the top left term. This inf-sup condition has been analytically established for various DPG methods through the construction of local [35, 11, 43] or global [12] Fortin operators on generously large test spaces. The same inf-sup condition also confirms the stability of the corresponding DPG* methods. An alternative characterization of the methods above can be found in a Petrov–Galerkin form in [39, Section 4.1].

Upon the choice of bases $\{v_i\}$ and $\{w_j\}$ for the discrete spaces \mathcal{V}_h and \mathcal{U}_h , (27) can be identified with the following system of matrix equations:

$$(28) \quad \begin{bmatrix} \mathbf{G} & \mathbf{B} \\ \mathbf{B}^T & 0 \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{w} \end{bmatrix} = \begin{bmatrix} \mathbf{f} \\ \mathbf{g} \end{bmatrix}.$$

Here, \mathbf{B} is a rectangular matrix with coefficients determined by the bilinear form, $B_{ij} = b(w_j, v_i)$, and, by conventional notation, \mathbf{G} is a Gram matrix governed by the chosen inner product, $G_{ik} = (v_i, v_k)_{\mathcal{V}}$. Naturally, the vectors $\mathbf{f}_i = F(v_i)$, $\mathbf{g}_j = G(w_j)$ are identified with the two loads in (27) and the vectors \mathbf{v} and \mathbf{w} correspond to the coefficients of the chosen basis functions. In the broken space setting (26), the Gram matrix can be block-diagonal. In that case, inverting

\mathbf{G} is computationally feasible and the Schur complement of (28) (cf. (9) and (11)) may be used to solve for the vector \mathbf{w} in a much smaller system, independent of \mathbf{v} :

$$(29) \quad \mathbf{B}^\top \mathbf{G}^{-1} \mathbf{B} \mathbf{w} = \mathbf{B}^\top \mathbf{G}^{-1} \mathbf{f} - \mathbf{g}.$$

Notice that the DPG stiffness matrix, $\mathbf{B}^\top \mathbf{G}^{-1} \mathbf{B}$, is always symmetric and positive-definite and that after solving for \mathbf{w} via (29), \mathbf{v} can always be recovered with only local cost, i.e., $\mathbf{v} = \mathbf{G}^{-1}(\mathbf{f} - \mathbf{B} \mathbf{w})$. Construction of the stiffness matrix $\mathbf{B}^\top \mathbf{G}^{-1} \mathbf{B}$ with broken spaces is considered in detail in [42].

2.3. Ultraweak formulations. Many PDEs originate in the following strong form:

$$(30) \quad \mathcal{L}u = f,$$

where \mathcal{L} is a linear differential operator and f is a prescribed function. It is possible to give many general DPG and DPG* formulations for such operator equations using the framework of [23, Appendix A] (which generalizes the Friedrichs systems framework in [27, 9, 53]). Let $d, k, m, l \geq 1$ be integers and let $\Omega \subseteq \mathbb{R}^d$ be a bounded open set. We use multiindices $\alpha = (\alpha_1, \dots, \alpha_d)$ of length $|\alpha| = \alpha_1 + \dots + \alpha_d \leq k$. Suppose we are given functions $a_{ij\alpha} : \Omega \rightarrow \mathbb{R}$ for each $i = 1, \dots, l, j = 1, \dots, m$, and each $|\alpha| \leq k$. Let \mathcal{L} be the differential operator acting on functions $u : \Omega \rightarrow \mathbb{R}^m$ such that

$$[\mathcal{L}u]_i = \sum_{j=1}^m \sum_{|\alpha| \leq k} \partial^\alpha (a_{ij\alpha} u_j), \quad i \in \{1, \dots, l\}.$$

Wherever appropriate, let L^2 denote *either* the l - or m -fold Cartesian product of $L^2(\Omega)$. Likewise, let \mathcal{D} denote *either* the l - or m -fold Cartesian product of $\mathcal{D}(\Omega)$, where $\mathcal{D}(\Omega)$ is the space of infinitely differentiable functions that are compactly supported on Ω (and accordingly, \mathcal{D}' denotes distributional vector fields). Let \mathcal{L}^* be the formal adjoint differential operator of \mathcal{L} , i.e., it satisfies $(\mathcal{L}\phi, \psi)_{L^2} = (\phi, \mathcal{L}^*\psi)_{L^2}$ for all $\phi, \psi \in \mathcal{D}$. From now on, we will simply denote all such L^2 -inner products on Ω as $(\cdot, \cdot)_\Omega = (\cdot, \cdot)_{L^2}$. Likewise, all L^2 -inner products restricted to a measurable subset $K \subseteq \Omega$ will be denoted $(\cdot, \cdot)_K$.

The action of \mathcal{L}^* on $v : \Omega \rightarrow \mathbb{R}^l$ is given by

$$(31) \quad [\mathcal{L}^*v]_j = \sum_{i=1}^l \sum_{|\alpha| \leq k} (-1)^{|\alpha|} a_{ij\alpha} \partial^\alpha v_i, \quad j \in \{1, \dots, m\}.$$

We assume that the coefficients $a_{ij\alpha}$ are such that both $\mathcal{L}u$ and \mathcal{L}^*v are well-defined distributions for all $u, v \in L^2$, i.e.,

$$(32a) \quad \mathcal{L}u \text{ and } \mathcal{L}^*v \text{ are in } \mathcal{D}' \text{ for all } u, v \in L^2.$$

(This holds e.g., if $a_{ij\alpha}$ are constant [23].)

We may now define Sobolev-like graph spaces by virtue of (32a). On any nonempty open subset $K \subseteq \Omega$, define the Hilbert spaces $H(\mathcal{L}, K) = \{u \in L^2(K)^m : \mathcal{L}u \in L^2(K)^l\}$ and, likewise, $H(\mathcal{L}^*, K) = \{v \in L^2(K)^l : \mathcal{L}^*v \in L^2(K)^m\}$. (E.g., if we let $\mathcal{L} = \text{grad}$, the canonical gradient operator, then $\mathcal{L}^* = -\text{div}$ and $H(\mathcal{L}, K) = H(\text{grad}, K) = H^1(K)$ and $H(\mathcal{L}^*, K) = H(\text{div}, K)$.) To simplify notation, we abbreviate $H(\mathcal{L}) = H(\mathcal{L}, \Omega)$ and $H(\mathcal{L}^*) = H(\mathcal{L}^*, \Omega)$. Also define linear operators $\mathcal{D} : H(\mathcal{L}) \rightarrow H(\mathcal{L}^*)'$ and $\mathcal{D}^* : H(\mathcal{L}^*) \rightarrow H(\mathcal{L})'$ such that

$$\langle \mathcal{D}u, v \rangle_{H(\mathcal{L}^*)} = (\mathcal{L}u, v)_\Omega - (u, \mathcal{L}^*v)_\Omega, \quad \langle \mathcal{D}^*v, u \rangle_{H(\mathcal{L})} = (\mathcal{L}^*v, u)_\Omega - (v, \mathcal{L}u)_\Omega,$$

for all $u \in H(\mathcal{L})$ and $v \in H(\mathcal{L}^*)$. Note that $\mathcal{D}^* = -\mathcal{D}'$, by these definitions. These graph spaces are equipped with natural *graph norms*:

$$\|u\|_{H(\mathcal{L})}^2 = \|\mathcal{L}u\|_{L^2}^2 + \|u\|_{L^2}^2, \quad \|v\|_{H(\mathcal{L}^*)}^2 = \|\mathcal{L}^*v\|_{L^2}^2 + \|v\|_{L^2}^2.$$

With these norms, notice that both \mathcal{D} and \mathcal{D}^* are bounded. Indeed, $|\langle \mathcal{D}u, v \rangle_{H(\mathcal{L}^*)}| \leq \|\mathcal{L}u\|_{L^2} \|v\|_{L^2} + \|u\|_{L^2} \|\mathcal{L}^*v\|_{L^2} \leq \|u\|_{H(\mathcal{L})} \|v\|_{H(\mathcal{L}^*)}$.

Finally, we may incorporate homogeneous boundary conditions. Recall the definition of the left annihilator in (7). Define $H_0(\mathcal{L}) \subseteq H(\mathcal{L})$ and $H_0(\mathcal{L}^*) \subseteq H(\mathcal{L}^*)$ to be two subspaces satisfying

$$(32b) \quad H_0(\mathcal{L}) = {}^\perp \mathcal{D}^*(H_0(\mathcal{L}^*)), \quad H_0(\mathcal{L}^*) = {}^\perp \mathcal{D}(H_0(\mathcal{L})).$$

Observe that (32b) does not uniquely characterize either $H_0(\mathcal{L})$ or $H_0(\mathcal{L}^*)$. These definitions permit many different so-called “mixed” homogeneous boundary conditions.

We will consider two boundary value problems: Given $f, g \in L^2$,

$$(33a) \quad \text{find } u \in H_0(\mathcal{L}) \text{ satisfying } \mathcal{L}u = f,$$

$$(33b) \quad \text{find } v \in H_0(\mathcal{L}^*) \text{ satisfying } \mathcal{L}^*v = g.$$

To derive a broken “ultraweak formulation” for (33a) and (33b), we focus on the scenario where Ω is partitioned into a mesh Ω_h of finitely many open disjoint elements K such that $\bar{\Omega}$ is the union of closures of all mesh elements K in Ω_h . For functions u and v , we denote by $\mathcal{L}_h u$ and $\mathcal{L}_h^* v$ the functions obtained by applying \mathcal{L} and \mathcal{L}^* to $u|_K$ and $v|_K$, respectively, element by element, for all $K \in \Omega_h$. With this in mind, define the broken spaces

$$H(\mathcal{L}_h) = \prod_{K \in \Omega_h} H(\mathcal{L}, K), \quad H(\mathcal{L}_h^*) = \prod_{K \in \Omega_h} H(\mathcal{L}^*, K),$$

which naturally conform to (26).

Clearly, $H(\mathcal{L}_h)$ and $H(\mathcal{L}_h^*)$ are inner product spaces with corresponding graph norms. The natural inner products on these spaces, induced by these graph norms, are defined

$$(34) \quad (u, \tilde{u})_{H(\mathcal{L}_h)} = (\mathcal{L}_h u, \mathcal{L}_h \tilde{u})_\Omega + (u, \tilde{u})_\Omega, \quad (v, \tilde{v})_{H(\mathcal{L}_h^*)} = (\mathcal{L}_h^* v, \mathcal{L}_h^* \tilde{v})_\Omega + (v, \tilde{v})_\Omega,$$

for all $u, \tilde{u} \in H(\mathcal{L}_h)$, $v, \tilde{v} \in H(\mathcal{L}_h^*)$. Now define the corresponding bounded linear operators $\mathcal{D}_h : H(\mathcal{L}_h) \rightarrow H(\mathcal{L}_h^*)'$ and $\mathcal{D}_h^* : H(\mathcal{L}_h^*) \rightarrow H(\mathcal{L}_h)'$ by

$$\langle \mathcal{D}_h u, v \rangle_{H(\mathcal{L}_h^*)} = (\mathcal{L}_h u, v)_\Omega - (u, \mathcal{L}_h^* v)_\Omega, \quad \langle \mathcal{D}_h^* v, u \rangle_{H(\mathcal{L}_h)} = (\mathcal{L}_h^* v, u)_\Omega - (v, \mathcal{L}_h u)_\Omega,$$

for all $u \in H(\mathcal{L}_h)$, $v \in H(\mathcal{L}_h^*)$. From now on, when using the operators \mathcal{D}_h and \mathcal{D}_h^* , we will simply denote $\langle \mathcal{D}_h \cdot, \cdot \rangle_h = \langle \mathcal{D}_h \cdot, \cdot \rangle_{H(\mathcal{L}_h^*)}$ or, likewise, $\langle \mathcal{D}_h^* \cdot, \cdot \rangle_h = \langle \mathcal{D}_h^* \cdot, \cdot \rangle_{H(\mathcal{L}_h)}$, since the meaning can easily be deduced from context. Finally, let

$$\begin{aligned} Q(\mathcal{L}_h) &= \{p \in H(\mathcal{L}_h)' : \text{there is a } v \in H_0(\mathcal{L}^*) \text{ such that } p = \mathcal{D}_h^* v\}, \\ Q(\mathcal{L}_h^*) &= \{q \in H(\mathcal{L}_h^*)' : \text{there is a } u \in H_0(\mathcal{L}) \text{ such that } q = \mathcal{D}_h u\}. \end{aligned}$$

These are Hilbert spaces when normed by the so-called *minimum energy extension* (quotient) norm [11, 36], i.e., $\|q\|_{Q(\mathcal{L}_h^*)} = \inf\{\|u\|_{H(\mathcal{L})} : u \in H(\mathcal{L}) \text{ satisfying } \mathcal{D}_h u = q\}$.

Multiplying (33a) by a function $\nu \in H(\mathcal{L}_h^*)$ and applying the definition of \mathcal{D}_h , we get $(u, \mathcal{L}_h^* \nu)_\Omega + \langle \mathcal{D}_h u, \nu \rangle_h = (f, \nu)_\Omega$ for all ν in $H(\mathcal{L}_h^*)$. Setting $\mathcal{D}_h u$ to q , a new unknown

in $Q(\mathcal{L}_h^*)$, we obtain the following *ultraweak formulation* with $F(\nu) = (f, \nu)_\Omega$. Given any $F \in H(\mathcal{L}_h^*)'$, find $u \in L^2$ and $q \in Q(\mathcal{L}_h^*)$ such that

$$(35a) \quad (u, \mathcal{L}_h^* \nu)_\Omega + \langle q, \nu \rangle_h = F(\nu) \quad \forall \nu \in H(\mathcal{L}_h^*).$$

Similarly proceeding with (33b) and setting $F(\nu) = (g, \nu)_\Omega$, we obtain an ultraweak formulation of the dual problem: Given any $F \in H(\mathcal{L}_h)'$, find $u \in L^2$ and $p \in Q(\mathcal{L}_h)$ such that

$$(35b) \quad (v, \mathcal{L}_h \nu)_\Omega + \langle p, \nu \rangle_h = F(\nu) \quad \forall \nu \in H(\mathcal{L}_h).$$

The next result shows that (35a) is uniquely solvable whenever (33a) is, as well as similar connection between (35b) and (33b).

Theorem 2.2 (Wellposedness of broken forms). *Suppose (32) holds. Then*

- (1) *Whenever $\mathcal{L} : H_0(\mathcal{L}) \rightarrow L^2$ is a bijection, problem (35a) is well-posed. Moreover, if $F(\nu) = (f, \nu)_\Omega$ for some $f \in L^2$, then the unique solution u of (35a) is in $H_0(\mathcal{L})$, solves (33a), and satisfies $q = \mathcal{D}_h u$.*
- (2) *Whenever $\mathcal{L}^* : H_0(\mathcal{L}^*) \rightarrow L^2$ is a bijection, problem (35b) is well-posed. Moreover, if $F(\nu) = (g, \nu)_\Omega$ for some $g \in L^2$, then the unique solution v of (35b) is in $H_0(\mathcal{L}^*)$, solves (33b), and satisfies $p = \mathcal{D}_h^* u$.*

Proof. The first statement is exactly the statement of [23, Theorem A.5]. The second statement also follows from [23, Theorem A.5] when \mathcal{L} is replaced by \mathcal{L}^* . \square

Naturally, formulations (35a) and (35b) also have adjoints. For instance, the adjoint of the ultraweak formulation (35a) is the following: Given any $G \in (L^2 \times Q(\mathcal{L}_h^*))'$, find $v \in H(\mathcal{L}_h^*)$ such that

$$(36a) \quad (\mu, \mathcal{L}_h^* v)_\Omega + \langle \rho, v \rangle_h = G(\mu, \rho) \quad \forall \mu \in L^2, \rho \in Q(\mathcal{L}_h^*).$$

Similarly, the adjoint of (35b) is: Given any $G \in (L^2 \times Q(\mathcal{L}_h))'$, find $u \in H(\mathcal{L}_h)$ such that

$$(36b) \quad (\mu, \mathcal{L}_h u)_\Omega + \langle \rho, u \rangle_h = G(\mu, \rho) \quad \forall \mu \in L^2, \rho \in Q(\mathcal{L}_h).$$

Under similar conditions to Theorem 2.2, these variational formulations are also well-posed, as the following theorem demonstrates.

Theorem 2.3 (Wellposedness of the adjoint problems). *Suppose (32) holds. Then*

- (1) *Whenever $\mathcal{L} : H_0(\mathcal{L}) \rightarrow L^2$ is a bijection, problem (36a) is well-posed. Moreover, if $G(\mu, \rho) = (g, \mu)_\Omega$ for some $g \in L^2$, then the unique solution v of (36a) is in $H_0(\mathcal{L}^*)$ and solves (33b).*
- (2) *Whenever $\mathcal{L}^* : H_0(\mathcal{L}^*) \rightarrow L^2$ is a bijection, problem (36b) is well-posed. Moreover, if $G(\mu, \rho) = (f, \mu)_\Omega$ for some $f \in L^2$, then the unique solution u of (36b) is in $H_0(\mathcal{L})$ and solves (33a).*

Proof. Both claims are closely related and follow similarly from Theorem 2.2. Therefore, we prove only the first statement.

Let the operator $\mathcal{B} : L^2 \times Q(\mathcal{L}_h^*) \rightarrow H(\mathcal{L}_h^*)'$ be defined $\langle \mathcal{B}(\mu, \rho), \nu \rangle_{H(\mathcal{L}_h^*)} = (\mu, \mathcal{L}_h^* \nu)_\Omega + \langle \rho, \nu \rangle_h$, for all $\nu \in H(\mathcal{L}_h)$ and $(\mu, \rho) \in L^2 \times Q(\mathcal{L}_h^*)$. Recall that $F \in H(\mathcal{L}_h^*)' = \text{Range } \mathcal{B}$ in (35a) was arbitrary. Therefore, as a consequence of the first statement in Theorem 2.2, we conclude that \mathcal{B} is both bounded below (cf. (5)) and surjective. That is, \mathcal{B} is a bijection and, by the Closed Range Theorem, $(\text{Null } \mathcal{B}')^\perp = \{0\}$. Hence, we conclude that (36a) is well-posed.

Next, suppose $G((\mu, \rho)) = (g, \mu)_\Omega$. Then (36a) yields

$$(37) \quad (\mu, \mathcal{L}_h^* v)_\Omega = (g, \mu)_\Omega,$$

$$(38) \quad \langle \rho, v \rangle_h = 0,$$

for all $\mu \in L^2$ and $\rho \in Q(\mathcal{L}_h^*)$. Equation (37) yields $\mathcal{L}_h^* v = g$ since $H(\mathcal{L}_h^*)$ is continuously embedded in L^2 . It remains to show that v is in $H_0(\mathcal{L}^*)$. Note that for all $\phi \in \mathcal{D}$, by the distributional definition of \mathcal{L} and the definition of \mathcal{D}_h ,

$$\langle \mathcal{L}^* v, \phi \rangle_{\mathcal{D}} = (\mathcal{L} \phi, v)_\Omega = (\mathcal{L}_h^* v, \phi)_\Omega + \langle \mathcal{D}_h \phi, v \rangle_h.$$

Since \mathcal{D} is contained in $H_0(\mathcal{L})$, $\mathcal{D}_h \phi$ is in $Q(\mathcal{L}_h^*)$ and the last term vanishes by virtue of (38). Moreover, since \mathcal{D} is densely contained in L^2 , this shows that $\mathcal{L}^* v = \mathcal{L}_h^* v = g$. Thus $v \in H(\mathcal{L}^*)$. Using (38) again, observe (cf. [23, Lemma A.3]) that

$$0 = \langle \rho, v \rangle_h = \langle \mathcal{D}_h \mu, v \rangle_h = \langle \mathcal{D} \mu, v \rangle_{H(\mathcal{L}^*)},$$

for all $\rho = \mathcal{D}_h \mu \in Q(\mathcal{L}_h^*)$, where $\mu \in H_0(\mathcal{L})$. Therefore, $v \in {}^\perp \mathcal{D}(H_0(\mathcal{L}))$. Finally, v is in $H_0(\mathcal{L}^*)$ simply by (32b). \square

Evidently, this result gives a class of examples where DPG* methods can be formulated. Letting $\mathcal{U} = L^2 \times Q(\mathcal{L}_h)$ and $\mathcal{V} = H_0(\mathcal{L}_h)$, define the bilinear form $b : \mathcal{U} \times \mathcal{V} \rightarrow \mathbb{R}$ as follows

$$(39) \quad b((\mu, \rho), \nu) = (\mu, \mathcal{L}_h \nu)_\Omega + \langle \rho, \nu \rangle_h \quad \forall (\mu, \rho) \in \mathcal{U}, \nu \in \mathcal{V}.$$

We may now consider the DPG* formulations of (33a). Treatment of the dual problem (33b) is similar.

Theorem 2.4 (Ultraweak DPG* formulation of (33a)). *Let $(\cdot, \cdot)_\mathcal{V}$ be any inner product on $H(\mathcal{L}_h)$ equivalent to $(\cdot, \cdot)_{H(\mathcal{L}_h)}$. Suppose (32) holds, $\mathcal{L}^* : H_0(\mathcal{L}^*) \rightarrow L^2$ is a bijection, and b is as in (39). Then, given a $G \in (L^2 \times Q(\mathcal{L}_h))'$, the problem of finding a function $u \in H(\mathcal{L}_h)$ satisfying*

$$(40) \quad \begin{cases} (u, \nu)_\mathcal{V} - b((\lambda, \sigma), \nu) &= 0 & \forall \nu \in H(\mathcal{L}_h), \\ b((\mu, \rho), u) &= G((\mu, \rho)) & \forall (\mu, \rho) \in L^2 \times Q(\mathcal{L}_h), \end{cases}$$

is well-posed. Moreover, if $G((\mu, \rho)) = (f, \mu)_\Omega$ for some $f \in L^2$, then the unique solution u is in $H_0(\mathcal{L})$ and satisfies $\mathcal{L}u = f$, i.e., u solves (33a).

Proof. Define the operator $B : H(\mathcal{L}_h) \rightarrow (L^2 \times Q(\mathcal{L}_h))'$, by $\langle B\nu, (\mu, \rho) \rangle_{L^2 \times Q(\mathcal{L}_h)} = b((\mu, \rho), \nu)$ for all $(\mu, \rho) \in L^2 \times Q(\mathcal{L}_h)$ and $\nu \in H(\mathcal{L}_h)$. As in the proof of Theorem 2.3, the operator $\mathcal{B} = B'$ is a bijection. Hence, (40) is a problem of form (4) (also (3)) with \mathcal{B} satisfying (5) and we conclude that (40) has a unique solution (see, e.g., Proposition 2.1). The remaining statements immediately follow from Theorem 2.3. \square

Remark 2.5. Note that the notation in Theorem 2.4 expresses the DPG* solution $u \in \mathcal{V}$ and $(\lambda, \sigma) \in \mathcal{U}$ in the form of (3). That is, the first solution component is denoted by the symbol u . From now on, in order to more closely follow the abstract notation used in (4), we will only use the symbol v for the \mathcal{V} -solution in all DPG* problems.

Remark 2.6. A very general broken space theory applicable of both DPG and DPG* methods has been established in the literature. This theory encompasses more traditional weak formulations and has been applied to a wide variety different boundary value problems [21, 8, 11, 41, 30]. For brevity, we will not expand on the intricate details here, but simply act to remind the

reader that ultraweak variational formulations are not a prerequisite for any DPG-type method coming from (27).

Example 2.7 (Poisson equation). In this example, which resurfaces throughout the document, $\vec{v} = (\vec{p}, v)$ will denote the DPG* solution variable. Similarly, $\vec{\lambda} = (\vec{\zeta}, \lambda, \hat{\zeta}_n, \hat{\lambda})$ will come to denote the associated Lagrange multiplier (see Section 3.2).

On a bounded open set $\Omega \subseteq \mathbb{R}^d$ with connected Lipschitz boundary, set $m = d + 1$ and

$$(41) \quad \mathcal{L}(\vec{p}, v) = (\vec{p} - \text{grad } v, -\text{div } \vec{p}),$$

where $\vec{p} : \Omega \rightarrow \mathbb{R}^d$ represents the flux variable and $v : \Omega \rightarrow \mathbb{R}$ represents the solution variable. Note that the equation $\mathcal{L}(\vec{p}, v) = (\vec{0}, f)$, after elimination of \vec{p} , results in the well-known Poisson equation $-\Delta v = f$.

We want to write out the DPG* formulation studied in Theorem 2.4 for this \mathcal{L} . Begin by observing that \mathcal{L}^* given by (31) can be written as

$$(42) \quad \mathcal{L}^*(\vec{\sigma}, \mu) = (\vec{\sigma} + \text{grad } \mu, \text{div } \vec{\sigma}).$$

Obviously (32a) is satisfied. In this case, by the triangle inequality, we immediately see that both $H(\mathcal{L})$ and $H(\mathcal{L}^*)$ coincide with $H(\text{div}, \Omega) \times H^1(\Omega)$. By integration by parts,

$$(43) \quad \langle \mathcal{D}^*(\vec{\sigma}, \mu), (\vec{p}, v) \rangle_h = \langle \vec{\sigma} \cdot \vec{n}, v \rangle_{H^{1/2}(\partial\Omega)} + \langle \vec{p} \cdot \vec{n}, \mu \rangle_{H^{1/2}(\partial\Omega)},$$

where \vec{n} denotes the unit outward normal on $\partial\Omega$. Put

$$(44) \quad H_0(\mathcal{L}^*) = H(\text{div}, \Omega) \times H_0^1(\Omega).$$

One may verify that this choice corresponds to the Dirichlet problem, $v = 0$ on $\partial\Omega$, and $H_0(\mathcal{L}) = H_0(\mathcal{L}^*)$.

From (43), it is immediate that (32b) holds. Along the lines of (43), we also have

$$\langle \mathcal{D}_h^*(\vec{\sigma}, \mu), (\vec{p}, v) \rangle_h = \sum_{K \in \Omega_h} \left[\langle \vec{\sigma} \cdot \vec{n}, v \rangle_{H^{1/2}(\partial K)} + \langle \vec{p} \cdot \vec{n}, \mu \rangle_{H^{1/2}(\partial K)} \right].$$

The range of $\mathcal{D}_h^*|_{H_0(\mathcal{L}^*)}$ is $Q(\mathcal{L}_h)$. In this example, this can be characterized using standard trace operators. The domain-dependent trace operators $\text{tr}^K u = u|_{\partial K}$ and $\text{tr}_n^K \vec{\sigma} = \vec{\sigma}|_{\partial K} \cdot \vec{n}$ for smooth functions are well-known to be continuously extendable to bounded linear maps $\text{tr}^K : H^1(K) \rightarrow H^{1/2}(\partial K)$ and $\text{tr}_n^K : H(\text{div}, K) \rightarrow H^{-1/2}(\partial K)$. Let $\text{tr} = \prod_{K \in \Omega_h} \text{tr}^K$ and $\text{tr}_n = \prod_{K \in \Omega_h} \text{tr}_n^K$. Then define

$$(45) \quad H^{-1/2}(\partial\Omega_h) = \text{tr}_n(H(\text{div}, \Omega)), \quad H_0^{1/2}(\partial\Omega_h) = \text{tr}(H_0^1(\Omega)).$$

Clearly, $Q(\mathcal{L}_h) = H^{-1/2}(\partial\Omega_h) \times H_0^{1/2}(\partial\Omega_h)$.

Applying the abstract setting to these definitions, the DPG* bilinear form in (39), becomes

$$(46) \quad \begin{aligned} b((\vec{\sigma}, \mu, \hat{\sigma}_n, \hat{\mu}), (\vec{\tau}, \nu)) &= \sum_{K \in \Omega_h} \left[(\vec{\sigma}, \vec{\tau} - \text{grad } \nu)_K - (\mu, \text{div } \vec{\tau})_K \right] \\ &+ \sum_{K \in \Omega_h} \left[\langle \vec{\tau} \cdot \vec{n}, \hat{\mu} \rangle_{H^{1/2}(\partial K)} + \langle \hat{\sigma}_n, \nu \rangle_{H^{1/2}(\partial K)} \right]. \end{aligned}$$

Here, $(\cdot, \cdot)_K$ denotes the inner product in $L^2(K)$ or its Cartesian products, $\vec{\sigma} \in L^2(\Omega)^d$, $\mu \in L^2(\Omega)$, $(\hat{\sigma}_n, \hat{\mu})$ is in the space $Q(\mathcal{L}_h)$ defined above, and the solution variable $(\vec{\sigma}, \mu)$ is in

the broken space $H(\mathcal{L}_h) = H(\operatorname{div}, \Omega_h) \times H^1(\Omega_h)$, where

$$H(\operatorname{div}, \Omega_h) = \prod_{K \in \Omega_h} H(\operatorname{div}, K), \quad H^1(\Omega_h) = \prod_{K \in \Omega_h} H^1(K).$$

Finally, the bijectivity of $\mathcal{L} : H_0(\mathcal{L}) \rightarrow L^2$ can be proved by standard techniques (see e.g. [19]). Hence Theorem 2.4 yields that this DPG* formulation is well posed.

We shall revisit this example later. In order to shorten the notation for later discussions, we shall denote the expression

$$(\operatorname{grad}_h \mu, \vec{\sigma})_\Omega + (\mu, \operatorname{div}_h \vec{\sigma})_\Omega$$

by $\langle \mu, \vec{\sigma} \cdot \vec{n} \rangle_h$ or $\langle \vec{\sigma} \cdot \vec{n}, \mu \rangle_h$ without explicitly indicating the application of the trace maps tr_n and tr . Accordingly, the bilinear form in (46) may be abbreviated as $b((\vec{\sigma}, \mu, \widehat{\sigma}_n, \widehat{\mu}), (\vec{\tau}, \nu)) = (\vec{\sigma}, \vec{\tau} - \operatorname{grad} \nu)_\Omega - (\mu, \operatorname{div} \vec{\tau})_\Omega + \langle \vec{\tau} \cdot \vec{n}, \widehat{\mu} \rangle_h + \langle \widehat{\sigma}_n, \nu \rangle_h$. \square

2.4. Related methods. Let $\mathcal{V} = H(\mathcal{L}_h^*)$. For any $F \in H(\mathcal{L}_h^*)'$, the ultraweak DPG formulation defined by (40) can be restated as the following system of variational equations:

$$(47a) \quad \begin{cases} (\varepsilon, \nu)_\mathcal{V} + (u, \mathcal{L}_h^* \nu)_\Omega + \langle p, \nu \rangle_h = F(\nu), & \forall \nu \in H(\mathcal{L}_h^*), \\ (\mu, \mathcal{L}_h^* \varepsilon)_\Omega = 0, & \forall \mu \in L^2, \\ \langle \rho, \varepsilon \rangle_h = 0, & \forall \rho \in Q(\mathcal{L}_h^*). \end{cases}$$

Likewise, letting $\mathcal{V} = H(\mathcal{L}_h)$, an ultraweak DPG formulation corresponding to (40) may be defined for any $G = G_\Omega \times G_h \in (L^2 \times Q(\mathcal{L}_h))'$:

$$(47b) \quad \begin{cases} (v, \nu)_\mathcal{V} - (\lambda, \mathcal{L}_h \nu)_\Omega - \langle \sigma, \nu \rangle_h = 0, & \forall \nu \in H(\mathcal{L}_h), \\ (\mu, \mathcal{L}_h v)_\Omega = G_\Omega(\mu), & \forall \mu \in L^2, \\ \langle \rho, v \rangle_h = G_h(\rho), & \forall \rho \in Q(\mathcal{L}_h). \end{cases}$$

Both of the formulations defined above relate to the primal problem (33a) with $u = v$. Clearly, the role of \mathcal{L}_h and \mathcal{L}_h^* can be interchanged if a solution of the dual problem (33b) is of interest.

The link between DPG and least-squares methods is well established in the literature (see e.g. [42]). DPG* methods, as it turns out, can be readily identified with the category of so-called \mathcal{LL}^* methods [10]. In this subsection, we briefly illustrate this and a couple of other notable relationships in the context of the mixed problems introduced in Section 2.1.

2.4.1. Least-squares methods. Let $\mathcal{V} = L^2$ and $\mathcal{U} = H_0(\mathcal{L})$. It is well-known that least-squares finite element methods [4] follow from the following saddle-point formulation (cf. (2) and (47a)):

$$(48) \quad \begin{cases} (\varepsilon, \nu)_\Omega + (\mathcal{L}u, \nu)_\Omega = F(\nu), & \forall \nu \in L^2, \\ (\mathcal{L}\mu, \varepsilon)_\Omega = 0, & \forall \mu \in H_0(\mathcal{L}). \end{cases}$$

This may be identified with a mixed problem akin to (2) using the strong formulation of (33a), rather than the ultraweak formulation, as in (47a). Indeed, let \mathcal{R}_{L^2} be the L^2 Riesz operator appearing in each term in (48) and recall identity (9), where $(\mathcal{B}\mu)(\cdot) = (\mathcal{L}\mu, \cdot)_\Omega$. Then observe that $\mathcal{B}' \mathcal{R}_\mathcal{V}^{-1} \mathcal{B} = (\mathcal{R}_{L^2} \mathcal{L})' \mathcal{R}_{L^2}^{-1} (\mathcal{R}_{L^2} \mathcal{L}) = \mathcal{L}' \mathcal{R}_{L^2} \mathcal{L}$ and $\mathcal{B}' \mathcal{R}_\mathcal{V}^{-1} F = \mathcal{L}' F$. That is,

$$\langle \mathcal{B}' \mathcal{R}_\mathcal{V}^{-1} \mathcal{B}u, \mu \rangle_\mathcal{U} = \langle \mathcal{B}' \mathcal{R}_\mathcal{V}^{-1} F, \mu \rangle_\mathcal{U} \iff (\mathcal{L}u, \mathcal{L}\mu)_\Omega = F(\mathcal{L}\mu).$$

In the case $F(\cdot) = (f, \cdot)_\Omega$, observe that $F(\mathcal{L}\nu) = (f, \mathcal{L}\nu)_\Omega$. Therefore, the variational equation above can be readily identified with the first-order optimality condition for the functional $\mathcal{J} : u \mapsto \|\mathcal{L}u - f\|_{L^2}^2$, $\partial_u \mathcal{J} = 0$.

2.4.2. \mathcal{LL}^* methods. Let $\mathcal{V} = L^2$ and $\mathcal{U} = H_0(\mathcal{L}^*)$. Contrary to (48), so-called \mathcal{LL}^* methods [10] relate to the following system (cf. (3) and (47b)):

$$\begin{cases} (v, \nu)_\Omega - (\mathcal{L}^* \lambda, \nu)_\Omega = 0, & \forall \nu \in L^2, \\ (\mathcal{L}^* \mu, v)_\Omega = G(\mu), & \forall \mu \in H_0(\mathcal{L}^*). \end{cases}$$

Likewise, consider (11), where $(\mathcal{B}\mu)(\cdot) = (\mathcal{L}^* \mu, \cdot)_\Omega$ and $G(\cdot) = (f, \cdot)_\Omega$. In this case, we see that \mathcal{LL}^* formulations may again be identified with (33a), in this case using a saddle-point expression akin to (3). Indeed, observe that

$$\langle \mathcal{B}' \mathcal{R}_\mathcal{V}^{-1} \mathcal{B} \lambda, \mu \rangle_\mathcal{U} = \langle G, \mu \rangle_\mathcal{U} \iff (\mathcal{L}^* \lambda, \mathcal{L}^* \mu)_\Omega = (f, \mu)_\Omega.$$

The variational equation above indicates, in a weak sense, that $\mathcal{LL}^* \lambda = f$. Recalling that the solution is determined by the transformation $v = \mathcal{R}_\mathcal{V}^{-1} \mathcal{B} \lambda = \mathcal{L}^* \lambda$, we have $\mathcal{L}v = f$ weakly, as well.

2.4.3. Weakly conforming least-squares methods. A weakly conforming least squares method [28] for the primal problem (33a) seeks a minimizer of the least squares functional

$$w \mapsto \|\mathcal{L}w - f\|_{L^2}^2,$$

under the conformity constraint

$$\langle w, \rho \rangle_h = 0, \quad \forall \rho \in Q(\mathcal{L}_h).$$

Here, of course, the operator \mathcal{L} is understood element-wise so we may saliently replace it by \mathcal{L}_h . This leads to the following saddle-point problem for the two solution components w and σ :

$$(49) \quad \begin{cases} (\mathcal{L}_h w, \mathcal{L}_h \nu)_\Omega + \langle \sigma, \nu \rangle_h = (f, \mathcal{L}_h \nu)_\Omega, & \forall \nu \in H(\mathcal{L}_h), \\ \langle w, \rho \rangle_h = 0, & \forall \rho \in Q(\mathcal{L}_h). \end{cases}$$

If we use an ultraweak DPG* formulation (47b) with its corresponding graph inner product (34), scaled by an arbitrary constant $\alpha > 0$, we arrive at

$$(50) \quad \begin{cases} (\mathcal{L}_h v, \mathcal{L}_h \nu)_\Omega + \alpha(v, \nu)_\Omega - (\lambda, \mathcal{L}_h \nu)_\Omega - \langle \sigma, \nu \rangle_h = 0, & \forall \nu \in H(\mathcal{L}_h), \\ (\mu, \mathcal{L}_h v)_\Omega = (f, \mu)_\Omega, & \forall \mu \in L^2, \\ \langle v, \rho \rangle_h = 0, & \forall \rho \in Q(\mathcal{L}_h). \end{cases}$$

From the second equation in (50), observe that $f = \mathcal{L}_h v$. Therefore, the first equation can be rewritten as

$$(f - \lambda, \mathcal{L}_h \nu)_\Omega + \alpha(v, \nu)_\Omega = \langle \sigma, \nu \rangle_h, \quad \forall \nu \in H(\mathcal{L}_h).$$

Testing only with $\nu \in H(\mathcal{L})$, so that the term $\langle \sigma, \nu \rangle_h$ vanishes, it can now be seen that $\lambda \rightarrow f$ as $\alpha \rightarrow 0$. Consequently, this particular DPG* formulation can be viewed as a regularization of the weakly conforming least-squares formulation (49).

2.5. Solving the primal and dual problems simultaneously. In (4), we may hypothetically consider any $F \in \mathcal{V}'$ and $G \in \mathcal{U}'$ we wish:

$$(51) \quad \begin{cases} \mathcal{R}_{\mathcal{V}} v + \mathcal{B} w = F, \\ \mathcal{B}' v = G. \end{cases}$$

Let \mathcal{B} be an isomorphism and define $F = \mathcal{R}_{\mathcal{V}}(\mathcal{B}')^{-1}G + \ell$, for some fixed $\ell \in (\text{Null } \mathcal{B}')^{\perp} = \mathcal{V}'$.³ Noting that $v = (\mathcal{B}')^{-1}G$, by the second equation in (51), it is readily seen that $\mathcal{B}w = \ell$. Therefore, with this choice of loads, $w = u$ solves the primal problem (1) and v solves the dual problem (10), simultaneously.

Introducing the load F , as proposed above, involves the inversion the linear operator \mathcal{B}' . In practice, this is usually not feasible and, therefore, precludes the construction of any such load in most circumstances. Nevertheless, consider the following system of equations:

$$(52) \quad \begin{cases} (\mathcal{L}^* v, \mathcal{L}^* \nu)_{\Omega} + (w, \mathcal{L}^* \nu)_{\Omega} = (g, \mathcal{L}^* \nu)_{\Omega} + (f, \nu)_{\Omega}, & \forall \nu \in H(\mathcal{L}^*), \\ (\mathcal{L}^* v, \mu)_{\Omega} = (g, \mu)_{\Omega}, & \forall \mu \in L^2. \end{cases}$$

This corresponds to a system like (51) with $\mathcal{V} = H(\mathcal{L}^*)$ and $\mathcal{U} = L^2$. In (52), v clearly satisfies $(\mathcal{L}^* v, \mu)_{\Omega} = (g, \mu)_{\Omega}$. That is, v solves the dual problem (33b), $\mathcal{L}^* v = g$, in a strong sense. Substituting $\mu = \mathcal{L}^* \nu$ into (52) and canceling terms in the first equation, we immediately find that $(w, \mathcal{L}^* \nu)_{\Omega} = (f, \nu)_{\Omega}$. That is, $w = u$ solves the primal problem (33a), $\mathcal{L}u = f$, in the ultraweak sense.

To avoid solving the mixed problem for both v and w at the same time, upon discretization, broken test spaces can be used. In this setting, we must consider the following related system with solution $(w, \sigma) \in \mathcal{U} = L^2 \times Q(\mathcal{L}_h^*)$ and $v \in \mathcal{V} = H(\mathcal{L}_h^*)$:

$$(53) \quad \begin{cases} (\mathcal{L}_h^* v, \mathcal{L}_h^* \nu)_{\Omega} + \alpha(v, \nu)_{\Omega} + (w, \mathcal{L}_h^* \nu)_{\Omega} + \langle \sigma, \nu \rangle_h = (g, \mathcal{L}_h^* \nu)_{\Omega} + (f, \nu)_{\Omega}, & \forall \nu \in H(\mathcal{L}_h^*), \\ (\mathcal{L}_h^* v, \mu)_{\Omega} = (g, \mu)_{\Omega}, & \forall \mu \in L^2, \\ \langle v, \rho \rangle_h = 0, & \forall \rho \in Q(\mathcal{L}_h^*). \end{cases}$$

Here the parameter $\alpha > 0$ has been added only to ensure that the $(1, 1)$ -block of the discrete system is locally invertible. Due to the constraint $\langle v, \rho \rangle_h = 0$, it can be shown that (53) is well-posed for any $\alpha \geq 0$.

The consequent manipulations are inspired by [33, Lemma 7]. First, notice that the last two equations in (53) uniquely determine v . Therefore, after testing the middle equation with $\mu = \mathcal{L}_h^* \nu$, observe that the first equation can be rewritten

$$(w, \mathcal{L}_h^* \nu)_{\Omega} + \langle \sigma, \nu \rangle_h = (f, \nu)_{\Omega} - \alpha(v, \nu).$$

By linearity, $(w, \sigma)_{\Omega} = (u, q)_{\Omega} + \alpha(e, r)$, where $(u, q) \in \mathcal{U}$ solves the ultraweak primal problem $(u, \mathcal{L}_h^* \nu)_{\Omega} + \langle q, \nu \rangle_h = (f, \nu)_{\Omega}$ (cf. (33a)) and $(e, r) = (e(v), r(v))$ is a *pollution term* defined by the equation $(e, \mathcal{L}_h^* \nu)_{\Omega} + \langle r, \nu \rangle_h = -(v, \nu)$. Clearly, $w \rightarrow u$ as $\alpha \rightarrow 0+$.

3. A PRIORI ERROR ANALYSIS

3.1. General results. Having explained the connections between the DPG* method and the mixed formulation (4), it should not be a surprise that its error analysis reduces to standard

³In the case of an injective but *not* surjective \mathcal{B} , consider $F = \mathcal{B}(\mathcal{B}' \mathcal{R}_{\mathcal{V}}^{-1} \mathcal{B})^{-1}G + \ell \in (\text{Null } \mathcal{B}')^{\perp} \subsetneq \mathcal{V}'$.

$$(54) \quad \begin{cases} (v, \nu)_{\mathcal{V}} - b(\lambda, \nu) = 0, & \nu \in \mathcal{V}, \\ b(\mu, v) = G(\mu), & \mu \in \mathcal{U}. \end{cases}$$
$$(55) \quad \begin{cases} (v_h, \nu)_{\mathcal{V}} - b(\lambda_h, \nu) = 0, & \nu \in \mathcal{V}_h, \\ b(\mu, v_h) = G(\mu), & \mu \in \mathcal{U}_h. \end{cases}$$
$$(56) \quad b(\mu, \nu - \Pi_h \nu) = 0, \quad \mu \in \mathcal{U}_h, \nu \in \mathcal{V}.$$

Theorem 3.1. *Suppose (5) and (56) hold. Then there is a constant C such that the complete DPG^{*} solution $(v, \lambda) \in \mathcal{V} \times \mathcal{U}$ satisfies the error estimate*

$$\|v - v_h\|_{\mathcal{V}} + \|\lambda - \lambda_h\|_{\mathcal{U}} \leq C \left[\inf_{\nu \in \mathcal{V}_h} \|v - \nu\|_{\mathcal{V}} + \inf_{\mu \in \mathcal{U}_h} \|\lambda - \mu\|_{\mathcal{U}} \right].$$

$$(57a) \quad (\varepsilon, \nu)_V + b(u, \nu) = F(\nu),$$

$$(57b) \quad b(\mu, \varepsilon) = 0,$$

$$(58) \quad \inf_{\mu \in \mathcal{U}_h} \|u - \mu\|_{\mathcal{U}} \leq c_0(h) \|F\|_{\mathcal{V}}.$$

Theorem 3.2. *Suppose (58) holds in addition to the assumptions of Theorem 3.1. Then there exists a positive function $c_0(h)$, which goes to 0 as $h \rightarrow 0$, such that the error in the DPG* solution component v_h satisfies*

$$F(v - v_h) \leq c_0(h) \|\mathcal{B}\| \|F\|_{\mathcal{V}'} \left[\inf_{\nu \in \mathcal{V}_h} \|v - \nu\|_{\mathcal{V}}^2 + \inf_{\mu \in \mathcal{U}_h} \|\lambda - \mu\|_{\mathcal{U}}^2 \right]^{1/2}.$$

$$\begin{aligned} F(v - v_h) &= (\varepsilon, v - v_h)_V + b(u, v - v_h) && \text{by (57a),} \\ &= b(u, v - v_h) && \text{since } \varepsilon = 0, \\ &= b(u - \mu, v - v_h) && \text{by (54) and (55),} \\ &\leq c_0(h) \|\mathcal{B}\| \|F\|_{V'} \|v - v_h\|_V && \text{by (58).} \end{aligned}$$

The proof is completed by applying [Theorem 3.1](#). \square

It is interesting to note that the duality argument for the DPG* method uses a DPG formulation: the system (57) is clearly a DPG formulation. Vice versa, the duality argument for DPG methods uses DPG* formulations, as can be seen from the duality arguments in [6, 32, 33]. Even though these references did not use the name “DPG*,” one can see DPG* formulations at work within their proofs.

At this point, an essential difficulty in DPG* methods (that was not present in DPG methods) becomes clear. Consider using a DPG* form $b(\cdot, \cdot)$ given by [Theorem 2.4](#) for solving the primal problem $\mathcal{L}v = f$. Then the error in v_h computed by the DPG* method not only depends on the regularity of the solution v , but also on the regularity of an extraneous Lagrange multiplier λ . This is evident from the best approximation error bounds appearing in [Theorems 3.1](#) and [3.2](#). The following example will clarify this observation further.

3.2. Application to the Poisson example. Given $f \in L^2(\Omega)$, consider approximating the Dirichlet solution v

$$(59) \quad -\Delta v = f \quad \text{in } \Omega, \quad v = 0 \quad \text{on } \partial\Omega,$$

by the DPG* method. We follow the setting of [Example 2.7](#). Accordingly, we set

$$\mathcal{U} = L^2(\Omega)^d \times L^2(\Omega) \times H^{-1/2}(\partial\Omega_h) \times H_0^{1/2}(\partial\Omega_h), \quad \mathcal{V} = H(\text{div}, \Omega_h) \times H^1(\Omega_h),$$

where $H^{-1/2}(\partial\Omega_h)$ and $H_0^{1/2}(\partial\Omega_h)$ are defined in (45). The DPG* formulation (54) of (59) characterizes two variables,

$$\vec{v} = (\vec{p}, v) \in \mathcal{V}, \quad \vec{\lambda} = (\vec{\zeta}, \lambda, \hat{\zeta}_n, \hat{\lambda}) \in \mathcal{U},$$

satisfying

$$(60a) \quad ((\vec{p}, v), (\vec{\tau}, \nu))_{\mathcal{V}} - b((\vec{\zeta}, \lambda, \hat{\zeta}_n, \hat{\lambda}), (\vec{\tau}, \nu)) = 0,$$

$$(60b) \quad b((\vec{\sigma}, \mu, \hat{\sigma}_n, \hat{\mu}), (\vec{p}, v)) = (f, \mu)_{\Omega},$$

for all $\vec{v} = (\vec{\tau}, \nu)$ in \mathcal{V} and all $\vec{\mu} = (\vec{\sigma}, \mu, \hat{\sigma}_n, \hat{\mu})$ in \mathcal{U} . Here, $b(\cdot, \cdot)$ as given by (46) and, as before, $(\cdot, \cdot)_{\Omega}$ denotes the inner product in $L^2(\Omega)$ (or its Cartesian products).

By [Theorem 2.4](#), \mathcal{B} is a bijection, so obviously (5) holds. Let Ω_h be a shape regular mesh of simplices and let $P_p(K)$ denote the space of polynomials of degree at most p on the simplex K . Define $P_p(\partial K) = \{\mu : \mu|_E \in P_p(E) \text{ for all codimension-one sub-simplices } E \text{ of } K\}$ and set

$$(61) \quad \begin{aligned} P_p(\Omega_h) &= \prod_{K \in \Omega_h} P_p(K), & P_p(\partial\Omega_h) &= \prod_{K \in \Omega_h} P_p(\partial K), \\ \tilde{P}_p(\partial\Omega_h) &= P_p(\partial\Omega_h) \cap \text{tr}(H_0^1(\Omega)), & \hat{P}_p(\partial\Omega_h) &= P_p(\partial\Omega_h) \cap \text{tr}_n(H(\text{div}, \Omega)). \end{aligned}$$

Clearly $\tilde{P}_p(\partial\Omega_h)$ is a subspace of $H_0^{1/2}(\partial\Omega_h)$ which consists of continuous single-valued functions on mesh interfaces. It is also obvious that the space $\hat{P}_p(\partial\Omega_h)$ is a subspace of $H^{-1/2}(\partial\Omega_h)$. Furthermore, it may be shown that every $\hat{\sigma}_n \in \hat{P}_p(\partial\Omega_h)$ has a corresponding $\hat{\sigma}$ in $\prod_{K \in \Omega_h} \tilde{x}P_p(K) + P_p(K)^d$ such that $\hat{\sigma}_n = \hat{\sigma} \cdot n$. In particular, the value of $\hat{\sigma}_n|_{\partial K^+}$ has the opposite sign of the value of $\hat{\sigma}_n|_{\partial K^-}$ at every point of a mesh interface $E = \partial K^+ \cap \partial K^-$ (so $\hat{\sigma}_n$ is not single-valued

on E). To remember this orientation dependence, we often write a $\hat{\sigma}_n$ in $\hat{P}(\Omega_h)$ as $\hat{\sigma} \cdot \vec{n}$. A Fortin operator satisfying (56) for the case

$$\begin{aligned}\mathcal{U}_h &= \{(\vec{\sigma}, \mu, \hat{\sigma}_n, \hat{\mu}) \in \mathcal{U} : \vec{\sigma} \in P_p(\Omega_h)^d, \mu \in P_p(\Omega_h), \hat{\sigma}_n \in \hat{P}_p(\partial\Omega_h), \hat{\mu} \in \tilde{P}_{p+1}(\partial\Omega_h)\} \\ \mathcal{V}_h &= \{(\vec{\tau}, \nu) \in \mathcal{V} : \vec{\tau} \in P_{p+d}(\Omega_h)^d, \nu \in P_{p+d}(\Omega_h)\}.\end{aligned}$$

was constructed in [35].

To understand the practical convergence rates, we must understand the regularity of $\vec{\lambda}$. One way to do this is to write down the boundary value problem that $\vec{\lambda}$ satisfies, as done in [6, 32]. An alternate technique can be seen in [33], which directly manipulates the variational equation (60a) using the information in (60b). We follow the latter approach in the next proof.

Proposition 3.3. *The solution components $\vec{\zeta}, \lambda, \hat{\zeta}_n, \hat{\lambda}$ of the system (60) can be characterized using the remaining solution components, \vec{p} and v , and the function f as*

$$(62) \quad \begin{aligned}\vec{\zeta} &= \vec{p} + \vec{r}, & \hat{\zeta}_n &= 2\vec{p} \cdot \vec{n} + \vec{r} \cdot \vec{n}, \\ \lambda &= f + e, & \hat{\lambda} &= e,\end{aligned}$$

where (\vec{r}, e) is in the space $H_0(\mathcal{L}^*)$ defined in (44) and satisfies the Dirichlet problem $\mathcal{L}^*(\vec{r}, e) = (\vec{0}, v + 2f)$ where \mathcal{L}^* is as in (42). Specifically, $e \in H_0^1(\Omega)$ satisfies $-\Delta e = v + 2f$ and $\vec{r} = -\text{grad } e$.

Proof. By Theorem 2.4, we know that (60b) implies that \vec{p}, v satisfies $\mathcal{L}(\vec{p}, v) = (\vec{0}, f)$, i.e.,

$$(63) \quad \vec{p} - \text{grad } v = \vec{0}, \quad -\text{div } \vec{p} = f.$$

Next, we manipulate the first term of (60a) as follows:

$$\begin{aligned}((\vec{p}, v), (\vec{\tau}, \nu))_{\mathcal{V}} &= (\vec{p}, \vec{\tau})_{\Omega} + (\text{div } \vec{p}, \text{div } \vec{\tau})_{\Omega} + (v, \nu)_{\Omega} + (\text{grad } v, \text{grad } \nu)_{\Omega} \\ &= (\vec{p}, \vec{\tau} - \text{grad } \nu)_{\Omega} + (\text{div } \vec{p}, \text{div } \vec{\tau})_{\Omega} + (v, \nu)_{\Omega} + 2(\text{grad } v, \text{grad } \nu)_{\Omega} \\ &= (\vec{p}, \vec{\tau} - \text{grad } \nu)_{\Omega} + (f, -\text{div } \vec{\tau})_{\Omega} + (v, \nu)_{\Omega} \\ &\quad + 2 \sum_{K \in \Omega_h} \left[\langle \vec{n} \cdot \vec{p}, \nu \rangle_{H^{1/2}(\partial K)} - (\text{div } \vec{p}, \nu)_K \right] \\ &= b((\vec{p}, f, 2\vec{p} \cdot \vec{n}, 0), (\vec{\tau}, \nu)) + (v + 2f, \nu)_{\Omega},\end{aligned}$$

where we have used (63) twice. Now, let $\vec{r} \in L^2(\Omega)^d, e \in L^2(\Omega)$ and $(\hat{r}_n, \hat{e}) \in Q(\mathcal{L}_h)$ satisfy

$$b((\vec{r}, e, \hat{r}_n, \hat{e}), (\vec{\tau}, \nu)) = (v + 2f, \nu)_{\Omega}$$

for all $(\vec{\tau}, \nu) \in H(\mathcal{L}_h) = \mathcal{V}$. This is a variational equation of the form (35a). Hence, by the first item of Theorem 2.2, \vec{r} and e are unique. Moreover, $(\vec{r}, e) \in H_0(\mathcal{L}^*)$ satisfies $\mathcal{L}^*(\vec{r}, e) = (0, v + 2f)$, and $\vec{r} \cdot \vec{n}|_{\partial K} = \hat{r}_n|_{\partial K}, e|_{\partial K} = \hat{e}|_{\partial K}$ on all mesh element boundaries. Thus,

$$((\vec{p}, v), (\vec{\tau}, \nu))_{\mathcal{V}} = b((\vec{p} + \vec{r}, f + e, (2\vec{p} + \vec{r}) \cdot \vec{n}, e), (\vec{\tau}, \nu)).$$

Comparing this with (60a), the result follows. \square

We may now apply Theorem 3.1 along with standard Bramble-Hilbert arguments (see [35, Corollary 3.6] for details) to obtain convergence rates dictated by the following corollary to Theorem 3.1.

Corollary 3.4. *Let $h = \max_{K \in \Omega_h} \text{diam}(K)$, $d = 2, 3$, and let the assumptions of [Theorem 3.1](#) and [Proposition 3.3](#) hold. Let $\vec{v}_h = (\vec{p}_h, v_h) \in \mathcal{V}_h$ and $\vec{\lambda}_h = (\vec{\zeta}_h, \lambda_h, \hat{\zeta}_{n,h}, \hat{\lambda}_h) \in \mathcal{U}_h$ be the DPG* solutions to (55), with $G(\vec{\mu}) = (f, \mu)$. Let $e \in H_0^1(\Omega)$ satisfy $-\Delta e = v + 2f$. Then*

$$(64) \quad \|\vec{v} - \vec{v}_h\|_{\mathcal{V}} + \|\vec{\lambda} - \vec{\lambda}_h\|_{\mathcal{U}} \leq Ch^s (\|v\|_{H^{s+2}(\Omega)} + \|e\|_{H^{s+2}(\Omega)}),$$

for all $1/2 < s \leq p+1$.

Proof. Note that the left-hand side of the inequalities in [Theorem 3.1](#) and [Corollary 3.4](#) coincide. Therefore, our proof proceeds by showing that $\inf_{\vec{v} \in \mathcal{V}_h} \|\vec{v} - \vec{v}\|_{\mathcal{V}}^2 + \inf_{\vec{\mu} \in \mathcal{U}_h} \|\vec{\lambda} - \vec{\mu}\|_{\mathcal{U}}^2$ is bounded from above by the right-hand side of (64).

Let $1/2 < s \leq p+1$ and notice that $\|\vec{v} - \vec{v}\|_{\mathcal{V}}^2 = \|\vec{p} - \vec{p}\|_{H(\text{div}, \Omega_h)}^2 + \|v - v\|_{H^1(\Omega_h)}^2$. Therefore, the following inequality, $\inf_{\vec{v} \in \mathcal{V}_h} \|\vec{v} - \vec{v}\|_{\mathcal{V}}^2 \leq Ch^s (\|\vec{p}\|_{H^{s+1}(\Omega)} + \|v\|_{H^{s+1}(\Omega)})$, is immediate upon substituting $\vec{v} = \prod_{K \in \Omega_h} (\Pi_{\text{div}}^K \vec{p}, \Pi_{\text{grad}}^K v)$, where $\Pi_{\text{div}}^K : H(\text{div}, K) \cap H^s(K) \rightarrow \vec{x}P_p(K) + P_p(K)^d$ and $\Pi_{\text{grad}}^K : H^{1+s}(K) \rightarrow P_{p+1}(K)$ are the local Raviart–Thomas and nodal interpolation operators, for each element $K \in \Omega_h$. Since $\vec{p} = \text{grad } v$, we see that $\inf_{\vec{v} \in \mathcal{V}_h} \|\vec{v} - \vec{v}\|_{\mathcal{V}}^2 \leq Ch^s \|v\|_{H^{s+2}(\Omega)}$.

To handle the term $\inf_{\vec{\mu} \in \mathcal{U}_h} \|\vec{\lambda} - \vec{\mu}\|_{\mathcal{U}}$, we remark that it is well known (cf. [24]) that there also exist global interpolants $\Pi_{\text{grad}} v \in H_0^1(\Omega)$, $\Pi_{\text{div}} \vec{p} \in H(\text{div}, \Omega)$, and $\Pi \lambda \in L^2(\Omega)$ such that $\Pi_{\text{grad}} v|_K \in P_{p+1}(K)$, $\Pi_{\text{div}} \vec{p}|_K \in \vec{x}P_p(K) + P_p(K)^d$, and $\Pi \lambda|_K \in P_p(K)$, for all $K \in \Omega$. Moreover, there exist constants C , depending on the polynomial degree p and the shape of the domain Ω , such that

$$(65a) \quad \|v - \Pi_{\text{grad}} v\|_{H^1(\Omega)} \leq Ch^r \|v\|_{H^{s+1}(\Omega)}, \quad (1/2 < r \leq p+1),$$

$$(65b) \quad \|\vec{p} - \Pi_{\text{div}} \vec{p}\|_{H(\text{div}, \Omega)} \leq Ch^r \|\vec{p}\|_{H^{s+1}(\Omega)}, \quad (0 < r \leq p+1),$$

$$(65c) \quad \|\lambda - \Pi \lambda\|_{L^2(\Omega)} \leq Ch^r \|\lambda\|_{H^s(\Omega)}, \quad (0 < r \leq p+1).$$

Notice that $\|\vec{\lambda} - \vec{\mu}\|_{\mathcal{U}}^2 = \|\vec{\zeta} - \vec{\sigma}\|_{L^2(\Omega)}^2 + \|\lambda - \mu\|_{L^2(\Omega)}^2 + \|\hat{\zeta}_n - \hat{\sigma}_n\|_{H^{-1/2}(\partial\Omega_h)}^2 + \|\hat{\lambda} - \hat{\mu}\|_{H^{1/2}(\partial\Omega_h)}^2$. Consider $\inf_{\vec{\sigma} \in L^2(\Omega)^d} \|\vec{\zeta} - \vec{\sigma}\|_{L^2(\Omega)} \leq Ch^s |\vec{\zeta}|_{H^s(\Omega)}$ and $\inf_{\mu \in L^2(\Omega)} \|\lambda - \mu\|_{L^2(\Omega)} \leq h^s |\lambda|_{H^s(\Omega)}$, both by (65c). Now, by (62) in [Proposition 3.3](#), $|\vec{\zeta}|_{H^s(\Omega)} \leq |\vec{p}|_{H^s(\Omega)} + |\vec{r}|_{H^s(\Omega)}$. Similarly, by invoking the identity $f = -\Delta v$, we have $|\lambda|_{H^s(\Omega)} \leq 2|f|_{H^s(\Omega)} + |e|_{H^s(\Omega)} \leq 2\|v\|_{H^{s+2}(\Omega)} + |e|_{H^s(\Omega)}$. Next, recall that $\|\text{tr}_n \vec{\sigma}\|_{H^{-1/2}(\partial\Omega_h)} \leq C \|\vec{\sigma}\|_{H(\text{div}, \Omega_h)}$, for any $\vec{\sigma} \in H(\text{div}, \Omega_h)$, by continuity of the normal trace operator $\text{tr}_n : H(\text{div}, \Omega_h) \rightarrow H^{-1/2}(\partial\Omega_h)$. Therefore, invoking (65b) and (62), we see that $\inf_{\hat{\sigma}_n \in H^{-1/2}(\partial\Omega_h)} \|\hat{\zeta}_n - \hat{\sigma}_n\|_{H^{-1/2}(\partial\Omega_h)} \leq Ch^s (2|\vec{p}|_{H^{1+s}(\Omega)} + |\vec{r}|_{H^{1+s}(\Omega)})$. Likewise, using the trace theorem, (65a), and (62), we find $\inf_{\hat{\mu} \in H^{1/2}(\partial\Omega_h)} \|\hat{\lambda} - \hat{\mu}\|_{H^{1/2}(\partial\Omega_h)} \leq Ch^s |e|_{H^{1+s}(\Omega)}$. Recalling that $\vec{p} = \text{grad } v$ and $\vec{r} = -\text{grad } e$ completes the proof. \square

The conclusion from [Corollary 3.4](#) is that even if the solution v has high regularity throughout the entire domain and up to the boundary, the convergence rate of the DPG* method is still controlled by a pollution variable e , which may happen to be less regular than v . Indeed, by elliptic regularity [29], e , which satisfies $-\Delta e = v + 2f$, will be at least as regular as v in the interior of the domain, but may not be as regular up to the boundary.

To illustrate how to get higher order convergence rates in weaker norms using duality, we want to apply [Theorem 3.2](#). To this end, we require sufficient regularity in the solution of the dual problem. Consider the case of full regularity, where, for any $g \in L^2(\Omega)$, the solution

$u \in H_0^1(\Omega)$ of the Dirichlet problem $-\Delta u = g$ satisfies

$$(66) \quad \|u\|_{H^2(\Omega)} \leq C\|g\|_{L^2(\Omega)}.$$

The inequality above is well known to hold on convex polygonal domains. In this case, we apply [Theorem 3.2](#) with $F \in \mathcal{V}'$ defined

$$(67) \quad F((\vec{\tau}, \nu)) = (v - v_h, \nu)_\Omega.$$

Note that F only sees the error in the first solution component of $\vec{v}_h = (\vec{p}_h, v_h)$ and that

$$(68) \quad \|F\|_{\mathcal{V}'} \leq \|v - v_h\|_{L^2(\Omega)}.$$

We now need to verify (58), so let us consider the present analog of (57), with the functional F in (67):

$$(69) \quad \begin{cases} (\vec{\varepsilon}, \vec{v})_{\mathcal{V}} + b(\vec{u}, \vec{v}) = F(\vec{v}), & \forall \vec{v} \in \mathcal{V}, \\ b(\vec{\mu}, \vec{\varepsilon}) = 0, & \forall \vec{\mu} \in \mathcal{U}. \end{cases}$$

First, observe that [Theorem 2.2](#) implies \mathcal{B} is a bijection, so (57b) implies that $\vec{\varepsilon} = 0$. Therefore (57a) reduces to finding $\vec{u} = (\vec{q}, u) \in \mathcal{U}$, where $b(\vec{u}, \vec{v}) = F(\vec{v})$ for all $\vec{v} \in \mathcal{V}$. This is an equation of the form (35b). Hence, the second item of [Theorem 2.2](#) implies that $\mathcal{L}^*\vec{u} = (\vec{0}, v - v_h)$; i.e., we may write $\vec{u} = (-\text{grad } u, u) \in H_0(\mathcal{L}^*) = H(\text{div}, \Omega) \times H_0^1(\Omega)$ such that $u \in H_0^1(\Omega)$ satisfies $-\Delta u = v - v_h$. Now, due to the full regularity estimate (66) applied to u , we have $\|u\|_{H^2(\Omega)} \leq C\|v - v_h\|_{L^2(\Omega)}$.

We may now invoke the complement of [Corollary 3.4](#), [33, Theorem 6]:

Theorem 3.5. *Let $p \in \mathbb{N}_0$, let \vec{u} be the solution to (69) for some arbitrary $F \in \mathcal{V}'$. Then there exists a constant C , depending only on p and the shape regularity of Ω_h , such that*

$$\inf_{\vec{\mu} \in \mathcal{U}_h} \|\vec{u} - \vec{\mu}\|_{\mathcal{U}} \leq Ch^{p+1} (\|u\|_{H^{p+2}(\Omega)} + \|\vec{q}\|_{H^{p+1}(\Omega_h)}).$$

Using the fact that $\vec{q} = -\text{grad } u$, we now have the estimate

$$\inf_{\vec{\mu} \in \mathcal{U}_h} \|\vec{u} - \vec{\mu}\|_{\mathcal{U}} \leq Ch\|u\|_{H^2(\Omega)} \leq Ch\|v - v_h\|_{L^2(\Omega)}.$$

Which is of the same form as (58). Finally, it is clear that assumption (58) holds with $c_0(h) = h$. Then, (68) and [Theorem 3.2](#) imply

$$\|v - v_h\|_{L^2(\Omega)}^2 = F(v - v_h) \leq Ch\|v - v_h\|_{L^2(\Omega)} \left[\inf_{\vec{v} \in \mathcal{V}_h} \|\vec{v} - \vec{v}\|_{\mathcal{V}}^2 + \inf_{\vec{\mu} \in \mathcal{U}_h} \|\vec{\lambda} - \vec{\mu}\|_{\mathcal{U}}^2 \right]^{1/2},$$

which provides one higher order of convergence in the L^2 norm for the solution component v_h .

Ultimately, the upshot of the entire *a priori* error analysis above is that poor *a priori* convergence rates are possible with this method, even for infinitely smooth solutions v , due to the Lagrange multiplier λ , which may not be as smooth (cf. [Section 5.3](#)). Thus, without an adaptive algorithm that helps one capture irregular solutions, the DPG* method is generally impractical for high-order methods. We therefore proceed by studying *a posteriori* error control.

4. A POSTERIORI ERROR CONTROL

In this section, we will present an abstract *a posteriori* error estimator valid for all ultraweak DPG* formulations (see [Section 2.3](#)). We then proceed to work out the example of the Poisson problem in full detail. Note that abstract ultraweak formulations encompass many physical models besides the Poisson example given above. Other important examples with similar functional settings include convection-dominated diffusion [\[25, 14\]](#), Stokes flow [\[48, 13\]](#), linear elasticity [\[7, 12\]](#), and acoustics [\[34, 44, 51\]](#).

4.1. Designing error estimators for general ultraweak DPG* formulations. Consider the general setting of [Section 2.3](#) and the broken ultraweak DPG* formulation which is proved to be well posed in [Theorem 2.4](#). Namely, with \mathcal{L} set to the general partial differential operator in [\(30\)](#), the problem of finding a $v \in H_0(\mathcal{L})$ satisfying $\mathcal{L}v = f$ is reformulated as [\(40\)](#), where

$$\langle \mathcal{B}(\mu, \rho), \nu \rangle_{\mathcal{V}} = b((\mu, \rho), \nu) = (\mu, \mathcal{L}_h \nu)_{\Omega} + \langle \rho, \nu \rangle_h,$$

for all $(\mu, \rho) \in \mathcal{U} = L^2 \times Q(\mathcal{L}_h)$ and $\nu \in \mathcal{V} = H(\mathcal{L}_h)$. The DPG* method produces an approximation to v using two finite-dimensional subspaces $\mathcal{U}_h \subseteq \mathcal{U}$ and $\mathcal{V}_h \subseteq \mathcal{V}$. Let

$$(70) \quad \eta(\nu) = \sup_{\rho \in Q(\mathcal{L}_h)} \frac{\langle \rho, \nu \rangle_h}{\|\rho\|_{Q(\mathcal{L}_h)}}, \quad \nu \in \mathcal{V}_h.$$

This quantity can usually be interpreted as a “jump term” in applications. The message of the next theorem is that, notwithstanding the generality of the operators considered, the design of *a posteriori* error estimators for all ultraweak DPG* formulations reduces to obtaining upper and lower bounds for η .

Theorem 4.1. *Consider the ultraweak DPG* formulation [\(40\)](#) with $G((\mu, \rho)) = (f, \mu)$, for some $f \in L^2$. Suppose [\(32\)](#) holds and $\mathcal{L}^* : H_0(\mathcal{L}^*) \rightarrow L^2$ is a bijection. Then, for any $v_h \in \mathcal{V}_h$ (not necessarily equal to the DPG* solution),*

$$(71) \quad \|\mathcal{B}\|^{-1} \|G - \mathcal{B}'v_h\|_{\mathcal{U}'} \leq \|v - v_h\|_{\mathcal{V}} \leq \|\mathcal{B}^{-1}\| \|G - \mathcal{B}'v_h\|_{\mathcal{U}'}$$

and, moreover,

$$(72) \quad \|G - \mathcal{B}'v_h\|_{\mathcal{U}'}^2 = \|\mathcal{L}_h v_h - f\|_{\Omega}^2 + \eta(v_h)^2.$$

Proof. In view of [\(40\)](#), $v - v_h$ satisfies

$$(73) \quad \begin{cases} R_{\mathcal{V}}(v - v_h) + \mathcal{B}((\lambda, \sigma)) = -R_{\mathcal{V}}v_h, \\ \mathcal{B}'(v - v_h) = G - \mathcal{B}'v_h. \end{cases}$$

By [Theorem 2.4](#), \mathcal{B} is a bijection. Hence, applying the identity [\(22\)](#) of [Proposition 2.1](#) to the system [\(73\)](#), we arrive at

$$\|v - v_h\|_{\mathcal{V}} = \|G - \mathcal{B}'v_h\|_{\mathcal{U}'} = \sup_{\mu \in \mathcal{U}} \frac{\langle G - \mathcal{B}'v_h, \mu \rangle_{\mathcal{U}}}{\|\mu\|_{\mathcal{U}}},$$

where the second equality comes directly from definition [\(15\)](#). Next, recall that the norms $\|\cdot\|_{\mathcal{U}}$ and $\|\cdot\|_{\mathcal{U}'}$ are equivalent. Indeed, $\|\mathcal{B}^{-1}\|^{-1} \|\mu\|_{\mathcal{U}} \leq \|\mu\|_{\mathcal{U}} \leq \|\mathcal{B}\| \|\mu\|_{\mathcal{U}}$, for all $\mu \in \mathcal{U}$. The first result, [\(71\)](#), now follows immediately.

To arrive at (72), simply observe that

$$\begin{aligned} \|G - \mathcal{B}'v_h\|_{\mathcal{U}'}^2 &= \sup_{\mu \in L^2, \rho \in Q(\mathcal{L}_h)} \frac{|(f, \mu)_\Omega - b((\mu, \rho), v_h)|^2}{\|(\mu, \rho)\|_{\mathcal{U}}^2} \\ &= \sup_{\mu \in L^2, \rho \in Q(\mathcal{L}_h)} \frac{|(f - \mathcal{L}_h v_h, \mu)_\Omega - \langle \rho, v_h \rangle_h|^2}{\|\mu\|_\Omega^2 + \|\rho\|_{Q(\mathcal{L}_h)}^2} \\ &= \sup_{\mu \in L^2} \frac{|(f - \mathcal{L}_h v_h, \mu)_\Omega|^2}{\|\mu\|_\Omega^2} + \sup_{\rho \in Q(\mathcal{L}_h)} \frac{|\langle \rho, v_h \rangle_h|^2}{\|\rho\|_{Q(\mathcal{L}_h)}^2} \end{aligned}$$

and the second result also follows. \square

4.2. A posteriori error analysis for the Poisson example. In this subsection, we develop a computable quantity that is equivalent to the function η defined above, for the example of the DPG* method for Poisson equation. We then provide a complete analysis of reliability and efficiency of the resulting error estimator.

Recall the variational formulation derived in Example 2.7 for Poisson's equation. Its bilinear form (see (46)) is

$$b((\vec{\sigma}, \mu, \hat{\sigma}_n, \hat{\mu}), (\vec{\tau}, \nu)) = ((\vec{\sigma}, \mu), \mathcal{L}_h(\vec{\tau}, \nu))_\Omega + \langle \hat{\mu}, \vec{\tau} \cdot \vec{n} \rangle_h + \langle \hat{\sigma}_n, \nu \rangle_h,$$

where $\mathcal{L}_h(\nu, \vec{\tau}) = (\vec{\tau} - \text{grad}_h \nu, -\text{div}_h \vec{\tau})$. In this subsection, we proceed by assuming, for simplicity, that $\Omega \subseteq \mathbb{R}^2$, and that Ω_h is a geometrically conforming triangular shape-regular mesh. Let \mathcal{E} denote the set of all mesh edges and let $\mathcal{E}_{\text{int}} \subseteq \mathcal{E}$ be the set of all interior edges of Ω_h . Let h_E be the length of any edge $E \in \mathcal{E}$. Any $E \in \mathcal{E}_{\text{int}}$ has two adjacent elements K^+ and K^- such that $E = \partial K^+ \cap \partial K^-$. Let

$$[\![\vec{\tau} \cdot \vec{n}]\!] = \vec{\tau}_{K^+} \cdot \vec{n}_{K^+} + \vec{\tau}_{K^-} \cdot \vec{n}_{K^-}, \quad [\![\vec{\tau} \cdot \vec{n}^\perp]\!] = \vec{n}_{K^+}^\perp \cdot \vec{\tau}_{K^+} + \vec{n}_{K^-}^\perp \cdot \vec{\tau}_{K^-}.$$

Here, \vec{n}_K^\perp is the tangential unit vector; i.e., if $\vec{n}_K = (n_1, n_2)$ then $\vec{n}_K^\perp = (-n_2, n_1)$. If $E \in \mathcal{E} \setminus \mathcal{E}_{\text{int}}$ is an exterior edge on the boundary of an element K , then with \vec{n} equal to the outward unit normal on $\partial\Omega$, we simply set $[\![\vec{\tau} \cdot \vec{n}]\!] = \vec{\tau}_K \cdot \vec{n}$ and $[\![\vec{\tau} \cdot \vec{n}^\perp]\!] = \vec{n}^\perp \cdot \vec{\tau}_K$. Similarly, for any scalar function ν that may be discontinuous across an interface $E \in \mathcal{E}_{\text{int}}$, we define

$$[\![\nu \vec{n}]\!] = \nu_{K^+} \vec{n}_{K^+} + \nu_{K^-} \vec{n}_{K^-}$$

and set $[\![\nu]\!] = \nu_K \vec{n}$ on boundary edges $E \in \mathcal{E} \setminus \mathcal{E}_{\text{int}}$.

The setting for the DPG* method for the Poisson equation (of Example 2.7), including its discrete spaces $\mathcal{U}_h, \mathcal{V}_h$, is described in Section 3.2. We continue with these settings in this subsection. Let h denote the maximum of h_E over all $E \in \mathcal{E}$. When mesh dependent quantities A and B satisfy $A \leq CB$, with a positive constant C independent of h , then we write $A \lesssim B$. When $A \lesssim B$ and $B \lesssim A$, then we write $A \approx B$. The main result of this subsection is the next theorem.

Theorem 4.2. *Under the above settings, suppose $(\vec{p}, v) \in \mathcal{V}$ and $(\vec{p}_h, v_h) \in \mathcal{V}_h$ are the exact solution and the discrete DPG* solution of the Laplace problem, respectively. Then*

$$\|(\vec{p}, v) - (\vec{p}_h, v_h)\|_{\mathcal{V}} \approx \eta_i(\vec{p}_h, v_h)$$

for $i \in \{1, 2\}$, where the computable error estimators η_i are defined by

$$\begin{aligned}\eta_1(\vec{p}_h, v_h)^2 &= \|\mathcal{L}(\vec{p}_h, v_h) - f\|_\Omega^2 + \sum_{E \in \mathcal{E}_{\text{int}}} h_E \|\llbracket \vec{p}_h \cdot \vec{n} \rrbracket\|_{L^2(E)}^2 + \sum_{E \in \mathcal{E}} h_E \|\llbracket v_h \vec{n} \rrbracket\|_{H^1(E)}^2, \\ \eta_2(\vec{p}_h, v_h)^2 &= \|\mathcal{L}(\vec{p}_h, v_h) - f\|_\Omega^2 + \sum_{E \in \mathcal{E}_{\text{int}}} h_E \|\llbracket \vec{p}_h \cdot \vec{n} \rrbracket\|_{L^2(E)}^2 + \sum_{E \in \mathcal{E}} h_E^{-1} \|\llbracket v_h \vec{n} \rrbracket\|_{L^2(E)}^2.\end{aligned}$$

The remainder of this subsection is devoted to proving this theorem. The main idea is to apply [Theorem 4.1](#), but some intermediate results need to be established first. To this end, recall that there exists a H^1 -norm minimum energy extension $\mathcal{E}_{\text{grad}}$ that provides a continuous right inverse of $\text{tr} : H^1(\Omega) \rightarrow H^{1/2}(\partial\Omega_h)$. Indeed, for all $\hat{w} \in H^{1/2}(\partial\Omega_h)$, $\mathcal{E}_{\text{grad}}$ is defined using the pre-image set $\text{tr}^{-1}\{\hat{w}\}$ simply as

$$\mathcal{E}_{\text{grad}}(\hat{w}) = \arg \min_{w \in \text{tr}^{-1}\{\hat{w}\}} \|w\|_{H^1(\Omega)}.$$

Similarly, for all $\hat{q}_n \in H^{-1/2}(\partial\Omega_h)$, the continuous right inverse of $\text{tr}_n : H(\text{div}, \Omega) \rightarrow H^{-1/2}(\partial\Omega_h)$ is defined

$$\mathcal{E}_{\text{div}}(\hat{q}_n) = \arg \min_{\vec{q} \in \text{tr}_n^{-1}\{\hat{q}_n\}} \|\vec{q}\|_{H(\text{div}, \Omega)}.$$

Clearly these operators can also be applied element by element.

Before establishing a number of lemmas, we pause to construct two helpful observations. First, for any $\hat{q}_n \in \hat{P}_p(\partial\Omega_h)$, let \hat{q}_E denote the function in $\hat{P}_p(\partial\Omega_h)$ that vanishes on all edges of \mathcal{E} , except on the edge E where it equals $\hat{q}|_E$. Observe that $\mathcal{E}_{\text{div}}(\hat{q}_E)$ is supported only on $\Omega_E = \bigcup\{K \in \Omega_h : \text{meas}(\partial K \cap E) \neq \emptyset\}$. Second, let b_E denote the edge bubble of E (i.e., the product of the barycentric coordinates of the endpoints of E) and define $\tilde{P}_{p+2}^0(\partial\Omega_h) = \{\mu \in \text{tr}(P_{p+2}(\Omega_h) \cap H_0^1(\Omega)) : \text{on any edge } E \in \mathcal{E}, \mu|_E = r_p b_E \text{ for some } r_p \in P_p(E)\}$. Likewise, for any $\hat{\mu} \in \tilde{P}_{p+2}^0(\partial\Omega_h)$, the function $\mathcal{E}_{\text{grad}}(\hat{\mu})$ is supported only on Ω_E .

Our first lemma may be thought of as an inf-sup condition involving the space of edge bubbles $\tilde{P}_{p+2}^0(\partial\Omega_h)$.

Lemma 4.3. *For any degree $p \geq 0$ and for any $\vec{q}_h \in P_p(\Omega_h)^2$,*

$$\sum_{E \in \mathcal{E}_{\text{int}}} h_E \|\llbracket \vec{q}_h \cdot \vec{n} \rrbracket\|_{L^2(E)}^2 \lesssim \sup_{\hat{\mu} \in \tilde{P}_{p+2}^0(\partial\Omega_h)} \frac{\langle \vec{q}_h \cdot \vec{n}, \hat{\mu} \rangle_h^2}{\|\mathcal{E}_{\text{grad}}(\hat{\mu})\|_{H^1(\Omega)}^2}.$$

Proof. We shall use the following two estimates that can be proved by scaling arguments using finite dimensionality (see e.g. [\[52\]](#)). For all $\hat{w} \in \text{tr}(P_p(\Omega_h) \cap H_0^1(\Omega))$,

$$(75) \quad \|b_E \hat{w}\|_{L^2(E)}^2 \leq \|\hat{w}\|_{L^2(E)}^2 \lesssim (b_E \hat{w}, \hat{w})_E,$$

$$(76) \quad |\mathcal{E}_{\text{grad}}(b_E \hat{w}_E)|_{H^1(\Omega_E)} \lesssim h_E^{-1/2} \|\hat{w}\|_{L^2(E)},$$

where $(\cdot, \cdot)_E$ denotes the inner product of $L^2(E)$ and \hat{w}_E is as defined above. For any $\vec{q}_h \in P_p(\Omega_h)$ with nontrivial $\llbracket \vec{q}_h \cdot \vec{n} \rrbracket_E$, we have

$$\begin{aligned}(77) \quad h_E \|\llbracket \vec{q}_h \cdot \vec{n} \rrbracket\|_{L^2(E)}^2 &\lesssim h_E (b_E \llbracket \vec{q}_h \cdot \vec{n} \rrbracket, \llbracket \vec{q}_h \cdot \vec{n} \rrbracket)_E && \text{by (75)} \\ &\lesssim \frac{(b_E \llbracket \vec{q}_h \cdot \vec{n} \rrbracket, \llbracket \vec{q}_h \cdot \vec{n} \rrbracket)_E}{|\mathcal{E}_{\text{grad}}(b_E \llbracket \vec{q}_h \cdot \vec{n} \rrbracket_E)|_{H^1(\Omega_E)}} h_E^{1/2} \|\llbracket \vec{q}_h \cdot \vec{n} \rrbracket_E\|_{L^2(E)} && \text{by (76),}\end{aligned}$$

which, after canceling $h_E^{1/2} \|[\vec{q}_h \cdot \vec{n}]\|_{L^2(E)}$ from both sides, provides a local version of the result we want to prove.

To get to the global estimate, we will accumulate the contributions of jumps across each edge. For this, it's useful to observe that for all $\hat{\mu} \in \tilde{P}_{p+2}^0(\partial\Omega_h)$, we have

$$(78) \quad |\mathcal{E}_{\text{grad}}(\hat{\mu})|_{H^1(\Omega)}^2 \lesssim \sum_{E \in \mathcal{E}_{\text{int}}} |\mathcal{E}_{\text{grad}}(\hat{\mu}_E)|_{H^1(\Omega_E)}^2.$$

This can be seen beginning from the linearity of $\mathcal{E}_{\text{grad}}$ and the fact that $\mathcal{E}_K = \{E \in \mathcal{E} : \text{meas}(E \cap \partial K) \neq 0\}$ has fixed finite cardinality:

$$\begin{aligned} |\mathcal{E}_{\text{grad}}(\hat{\mu})|_{H^1(\Omega)}^2 &= \left| \sum_{E \in \mathcal{E}_{\text{int}}} \mathcal{E}_{\text{grad}}(\hat{\mu}_E) \right|_{H^1(\Omega)}^2 = \sum_{K \in \Omega_h} \left| \sum_{E \in \mathcal{E}_{\text{int}}} \mathcal{E}_{\text{grad}}(\hat{\mu}_E) \right|_{H^1(K)}^2 \\ &= \sum_{K \in \Omega_h} \left| \sum_{E \in \mathcal{E}_K} \mathcal{E}_{\text{grad}}(\hat{\mu}_E) \right|_{H^1(K)}^2 \lesssim \sum_{K \in \Omega_h} \sum_{E \in \mathcal{E}_K} |\mathcal{E}_{\text{grad}}(\hat{\mu}_E)|_{H^1(K)}^2 \lesssim \sum_{E \in \mathcal{E}_{\text{int}}} |\mathcal{E}_{\text{grad}}(\hat{\mu}_E)|_{H^1(\Omega_E)}^2, \end{aligned}$$

which proves (78).

We can now complete the proof as follows. Starting from (77),

$$\begin{aligned} \sum_{E \in \mathcal{E}_{\text{int}}} h_E \left\| [\vec{q}_h \cdot \vec{n}] \right\|_{L^2(E)}^2 &\lesssim \sum_{E \in \mathcal{E}_{\text{int}}} \frac{(b_E [\vec{q}_h \cdot \vec{n}], [\vec{q}_h \cdot \vec{n}])_E^2}{|\mathcal{E}_{\text{grad}}(b_E [\vec{q}_h \cdot \vec{n}])|_{H^1(\Omega_E)}^2} \\ &\leq \sum_{E \in \mathcal{E}_{\text{int}}} \sup_{\hat{\mu} \in \tilde{P}_{p+2}^0(\partial\Omega_h)} \frac{(\hat{\mu}_E, [\vec{q}_h \cdot \vec{n}])_E^2}{|\mathcal{E}_{\text{grad}}(\hat{\mu}_E)|_{H^1(\Omega_E)}^2} = \sup_{\hat{\mu} \in \tilde{P}_{p+2}^0(\partial\Omega_h)} \frac{\left(\sum_{E \in \mathcal{E}_{\text{int}}} (\hat{\mu}_E, [\vec{q}_h \cdot \vec{n}])_E \right)^2}{\sum_{E \in \mathcal{E}_{\text{int}}} |\mathcal{E}_{\text{grad}}(\hat{\mu}_E)|_{H^1(\Omega_E)}^2}, \end{aligned}$$

where, in the equality, we have exploited a property of suprema over components of a Cartesian product space (noting that the space $\tilde{P}_{p+2}^0(\partial\Omega_h)$ is the Cartesian product of $b_E P_p(E)$ over all interior edges E). Now, the result follows by noting that the numerator above equals $\langle \mu, \vec{q}_h \cdot \vec{n} \rangle_h^2$ and by bounding the denominator using (78) and the Poincaré inequality. \square

Lemma 4.4. *For any degree $p \geq 1$ and for any $w_h \in P_p(\Omega_h)$,*

$$\sum_{E \in \mathcal{E}_{\text{int}}} h_E \left\| [w_h \vec{n}] \right\|_{H^1(E)}^2 \lesssim \sup_{\hat{\sigma} \cdot \vec{n} \in \hat{P}_p(\partial\Omega_h)} \frac{\langle \hat{\sigma} \cdot \vec{n}, w_h \rangle_h^2}{\| \mathcal{E}_{\text{div}}(\hat{\sigma} \cdot \vec{n}) \|_{H(\text{div}, \Omega)}^2},$$

where $\hat{P}_p(\partial\Omega_h)$ is as defined in (61).

Proof. For any $w_h \in P_p(\Omega_h)$ and any $E \in \mathcal{E}$, the function $[\vec{n}^\perp \cdot \text{grad } w_h]$ represents the tangential derivative of the jump of w_h across E . Then $\phi_E = \mathcal{E}_{\text{grad}}([\vec{n}^\perp \cdot \text{grad } w_h]_E b_E)$ is supported on Ω_E and the trace of ϕ_E vanishes on all edges except E . Let $\Omega_{h,E} = \{K \in \Omega_h : K \subseteq \Omega_E\}$. Using the vector curl of the scalar function ϕ_E , by an application of (75), we have

$$\begin{aligned} \left\| [w_h \vec{n}] \right\|_{H^1(E)}^2 &\lesssim (b_E [\vec{n}^\perp \cdot \text{grad } w_h], [\vec{n}^\perp \cdot \text{grad } w_h])_E \\ &= \int_E \phi_E [\vec{n}^\perp \cdot \text{grad } w_h] = \sum_{K \in \Omega_{h,E}} \int_{\partial K} \phi_E \vec{n}^\perp \cdot \text{grad } w_h \\ &= \sum_{K \in \Omega_{h,E}} (\text{curl } \phi_E, \text{grad } w_h)_K = \sum_{K \in \Omega_{h,E}} \int_{\partial K} \vec{n} \cdot \text{curl } \phi_E w_h = (\text{curl } \phi_E, [w_h \vec{n}])_E, \end{aligned}$$

where we have also used the Stokes and the divergence theorems in succession. Now, noting that $\|\operatorname{curl} \phi_E\|_{H(\operatorname{div}, \Omega_E)} = |\phi_E|_{H^1(\Omega_E)} = |\mathcal{E}_{\operatorname{grad}}(\llbracket \operatorname{grad} w_h \cdot \vec{n}^\perp \rrbracket_E b_E)|_{H^1(\Omega_E)}$ we deduce using (76) that $\|\operatorname{curl} \phi_E\|_{H(\operatorname{div}, \Omega_E)} \lesssim h_E^{-1/2} \|\llbracket w_h \vec{n} \rrbracket\|_{H^1(E)}$. Hence

$$\|\llbracket w_h \vec{n} \rrbracket\|_{H^1(E)}^2 \lesssim \frac{(\operatorname{curl} \phi_E, \llbracket w_h \vec{n} \rrbracket)_E}{\|\operatorname{curl} \phi_E\|_{H(\operatorname{div}, \Omega_E)}} \|\operatorname{curl} \phi_E\|_{H(\operatorname{div}, \Omega_E)} \lesssim \frac{(\operatorname{curl} \phi_E, \llbracket w_h \vec{n} \rrbracket)_E}{\|\operatorname{curl} \phi_E\|_{H(\operatorname{div}, \Omega_E)}} h_E^{-1/2} \|\llbracket w_h \vec{n} \rrbracket\|_{H^1(E)}.$$

Together with the minimal extension property $\|\mathcal{E}_{\operatorname{div}}(\vec{n} \cdot \operatorname{curl} \phi_E)\|_{H(\operatorname{div}, \Omega)} \leq \|\operatorname{curl} \phi_E\|_{H(\operatorname{div}, \Omega)}$, this implies that

$$h_E \|\llbracket w_h \vec{n} \rrbracket\|_{H^1(E)}^2 \lesssim \frac{(\operatorname{curl} \phi_E, \llbracket w_h \vec{n} \rrbracket)_E}{\|\mathcal{E}_{\operatorname{div}}(\vec{n} \cdot \operatorname{curl} \phi_E)\|_{H(\operatorname{div}, \Omega_E)}^2} \leq \sup_{\widehat{\sigma}_E \cdot \vec{n} \in \widehat{P}_p(E)} \frac{\langle \widehat{\sigma}_E \cdot \vec{n}, w_h \rangle_h}{\|\mathcal{E}_{\operatorname{div}}(\widehat{\sigma}_E \cdot \vec{n})\|_{H(\operatorname{div}, \Omega_E)}^2},$$

where $\widehat{P}_p(E)$ denotes the subspace of functions in $\widehat{P}_p(\Omega_h)$ supported on E . We have thus arrived at a local version of the desired inequality. By proving an analogue of (78) for $\mathcal{E}_{\operatorname{div}}$, and following along the lines of the proof of Lemma 4.3, we finish the proof. \square

Since the suprema in Lemmas 4.3 and 4.4 are related to the function η in (70), these lemmas can be thought of providing lower bounds, often called *efficiency* estimates in the analysis of *a posteriori* estimators. To prove upper bounds, also called *reliability* estimates, we need some additional tools. Recall that any $\vec{\sigma} \in H(\operatorname{div}, \Omega)$ may be decomposed using the so-called *regular decomposition* [26, Section 3.3] as $\vec{\sigma} = \operatorname{curl}(\varphi_{\vec{\sigma}}) + \vec{\psi}_{\vec{\sigma}}$ such that

$$\|\varphi_{\vec{\sigma}}\|_{H^1(\Omega)} + \|\vec{\psi}_{\vec{\sigma}}\|_{H^1(\Omega)} \lesssim \|\vec{\sigma}\|_{H(\operatorname{div}, \Omega)}.$$

We shall also need low-regularity *commuting quasi-interpolators* of [26], built using refinements of earlier ideas in [15, 49]. Namely, there exist operators $\mathcal{I}_{\operatorname{grad}} : H^1(\Omega) \rightarrow P_1(\Omega_h) \cap H^1(\Omega)$ and $\mathcal{I}_{\operatorname{div}} : H(\operatorname{div}, \Omega) \rightarrow \mathcal{RT}_0(\Omega_h) \cap H(\operatorname{div}, \Omega)$, such that $\operatorname{curl} \circ \mathcal{I}_{\operatorname{grad}} = \mathcal{I}_{\operatorname{div}} \circ \operatorname{curl}$. Here, $\mathcal{RT}_0(\Omega_h) = P_0(\Omega_h)^2 + \vec{x}P_0(\Omega_h)$, the lowest order Raviart–Thomas space. In addition, the following inequalities [26, Lemma 6] hold for all $\mu \in H^1(\Omega)$ and $\vec{\sigma} \in H(\operatorname{div}, \Omega)$:

$$(79a) \quad \sum_{E \in \mathcal{E}} h_E^{-1} \|\mu - \mathcal{I}_{\operatorname{grad}} \mu\|_{L^2(E)}^2 \lesssim \|\mu\|_{H^1(\Omega)}^2,$$

$$(79b) \quad \sum_{E \in \mathcal{E}} h_E^{-1} \|\varphi_{\vec{\sigma}} - \mathcal{I}_{\operatorname{grad}} \varphi_{\vec{\sigma}}\|_{L^2(E)}^2 + h_E^{-1} \|(\vec{\psi}_{\vec{\sigma}} - \mathcal{I}_{\operatorname{div}} \vec{\psi}_{\vec{\sigma}}) \cdot \vec{n}\|_{L^2(E)}^2 \lesssim \|\vec{\sigma}\|_{H(\operatorname{div}, \Omega)}^2.$$

These results also hold with $H^1(\Omega)$ replaced by $H_0^1(\Omega)$ and $H(\operatorname{div}, \Omega)$ replaced by $H_0(\operatorname{div}, \Omega)$.

Lemma 4.5. *For any degree $p \geq 0$ and any $\vec{q}_h \in P_p(\Omega_h)^2$ satisfying $\langle \widehat{\mu}_1, \vec{q}_h \cdot \vec{n} \rangle_h = 0$ for all $\widehat{\mu}_1 \in \operatorname{tr}(P_1(\Omega_h) \cap H_0^1(\Omega))$, we have*

$$(80) \quad \sum_{E \in \mathcal{E}_{\operatorname{int}}} h_E \|\llbracket \vec{q}_h \cdot \vec{n} \rrbracket\|_{L^2(E)}^2 \approx \sup_{\mu \in H_0^1(\Omega)} \frac{\langle \mu, \vec{q}_h \cdot \vec{n} \rangle_h^2}{\|\mu\|_{H^1(\Omega)}^2}.$$

Proof. By Lemma 4.3,

$$\begin{aligned} \sum_{E \in \mathcal{E}_{\operatorname{int}}} h_E \|\llbracket \vec{q}_h \cdot \vec{n} \rrbracket\|_{L^2(E)}^2 &\lesssim \sup_{\widehat{\mu}_h \in \widetilde{P}_{p+2}^0(\partial\Omega_h)} \frac{\langle \vec{q}_h \cdot \vec{n}, \widehat{\mu}_h \rangle_h^2}{\|\mathcal{E}_{\operatorname{grad}}(\widehat{\mu}_h)\|_{H^1(\Omega)}^2} \lesssim \sup_{\widehat{\mu} \in H_0^{1/2}(\partial\Omega_h)} \frac{\langle \widehat{\mu}, \vec{q}_h \cdot \vec{n} \rangle_h^2}{\|\widehat{\mu}\|_{H^{1/2}(\partial\Omega_h)}^2} \\ &= \sup_{\mu \in H_0^1(\Omega)} \frac{\langle \mu, \vec{q}_h \cdot \vec{n} \rangle_h^2}{\|\mu\|_{H^1(\Omega)}^2}, \end{aligned}$$

where the last identity follows from previous works (see [19] or [11, Theorem 2.3]).

Hence, it suffices to prove the reverse inequality. Since $\langle \mathcal{I}_{\text{grad}} \mu, \vec{q}_h \cdot \vec{n} \rangle_h = 0$,

$$\begin{aligned} \langle \mu, \vec{q}_h \cdot \vec{n} \rangle_h &= \langle \mu - \mathcal{I}_{\text{grad}} \mu, \vec{q}_h \cdot \vec{n} \rangle_h = \sum_{E \in \mathcal{E}_{\text{int}}} (\mu - \mathcal{I}_{\text{grad}} \mu, \llbracket \vec{q}_h \cdot \vec{n} \rrbracket)_E \\ &\lesssim \sum_{E \in \mathcal{E}_{\text{int}}} h_E^{-1/2} \|\mu - \mathcal{I}_{\text{grad}} \mu\|_{L^2(E)} h_E^{1/2} \left\| \llbracket \vec{q}_h \cdot \vec{n} \rrbracket \right\|_{L^2(E)} \lesssim \|\mu\|_{H^1(\Omega)} \left(\sum_{E \in \mathcal{E}_{\text{int}}} h_E \left\| \llbracket \vec{q}_h \cdot \vec{n} \rrbracket \right\|_{L^2(E)}^2 \right)^{1/2}. \end{aligned}$$

Here, in the final line, we have used the Cauchy–Schwarz inequality and (79a). This completes the proof of (80). \square

Lemma 4.6. *For any degree $p \geq 1$ and any $w_h \in P_p(\Omega_h)$ satisfying $\int_E \llbracket w_h \vec{n} \rrbracket = 0$ on all edges $E \in \mathcal{E}$,*

$$(81) \quad \sum_{E \in \mathcal{E}} h_E \left\| \llbracket w_h \vec{n} \rrbracket \right\|_{H^1(E)}^2 \approx \sum_{E \in \mathcal{E}} h_E^{-1} \left\| \llbracket w_h \vec{n} \rrbracket \right\|_{L^2(E)}^2 \approx \sup_{\vec{\sigma} \in H(\text{div}, \Omega)} \frac{\langle \vec{\sigma} \cdot \vec{n}, w_h \rangle_h^2}{\|\vec{\sigma}\|_{H(\text{div}, \Omega)}^2}.$$

Proof. The first equivalence in (81) immediately follows from the Poincaré inequality since the mean value of $\llbracket w_h \vec{n} \rrbracket$ vanishes. To prove the remaining equivalence, first observe that Lemma 4.4 implies

$$\begin{aligned} \sum_{E \in \mathcal{E}_{\text{int}}} h_E \left\| \llbracket w_h \vec{n} \rrbracket \right\|_{H^1(E)}^2 &\lesssim \sup_{\hat{\sigma} \cdot \vec{n} \in \hat{P}_p(\partial\Omega_h)} \frac{\langle \hat{\sigma} \cdot \vec{n}, w_h \rangle_h^2}{\|\mathcal{E}_{\text{div}}(\hat{\sigma} \cdot \vec{n})\|_{H(\text{div}, \Omega)}^2} \\ &\leq \sup_{\hat{\sigma} \cdot \vec{n} \in H^{-1/2}(\partial\Omega_h)} \frac{\langle \hat{\sigma} \cdot \vec{n}, w_h \rangle_h^2}{\|\hat{\sigma} \cdot \vec{n}\|_{H^{-1/2}(\partial\Omega_h)}^2} = \sup_{\sigma \in H(\text{div}, \Omega)} \frac{\langle \sigma \cdot \vec{n}, w_h \rangle_h^2}{\|\sigma\|_{H(\text{div}, \Omega)}^2}, \end{aligned}$$

where the last identity is well known (see [11, Theorem 2.3]). This proves one side of the stated equivalence.

To prove the remaining inequality, we start by decomposing any given $\vec{\sigma} \in H(\text{div}, \Omega)$ using above-mentioned regular decomposition: $\vec{\sigma} = \text{curl } \varphi_{\vec{\sigma}} + \vec{\psi}_{\vec{\sigma}}$. Then, since the jump of w_h has zero mean value on every edge, we observe that $\langle \vec{n} \cdot \mathcal{I}_{\text{div}} \vec{\psi}_{\vec{\sigma}}, w_h \rangle_h = \langle \vec{n} \cdot \mathcal{I}_{\text{div}}(\text{curl } \varphi_{\vec{\sigma}}), w_h \rangle_h = 0$. Therefore, by the commutativity property of the quasi-interpolators,

$$\langle \vec{\sigma} \cdot \vec{n}, w_h \rangle_h = \langle \vec{n} \cdot (\text{curl } \varphi_{\vec{\sigma}} - \mathcal{I}_{\text{grad}} \varphi_{\vec{\sigma}}), w_h \rangle_h + \langle \vec{n} \cdot (\vec{\psi}_{\vec{\sigma}} - \mathcal{I}_{\text{div}} \vec{\psi}_{\vec{\sigma}}), w_h \rangle_h.$$

We proceed labeling the terms on the right as t_1 and t_2 , respectively. Using the divergence theorem and the Stokes theorem in succession,

$$\begin{aligned} t_1 &= (\text{curl}(\varphi_{\vec{\sigma}} - \mathcal{I}_{\text{grad}} \varphi_{\vec{\sigma}}), \text{grad } w_h)_\Omega \\ &= \langle \varphi_{\vec{\sigma}} - \mathcal{I}_{\text{grad}} \varphi_{\vec{\sigma}}, \vec{n}^\perp \cdot \text{grad } w_h \rangle_h = \sum_{E \in \mathcal{E}} (\varphi_{\vec{\sigma}} - \mathcal{I}_{\text{grad}} \varphi_{\vec{\sigma}}, \llbracket \vec{n}^\perp \cdot \text{grad } w_h \rrbracket)_E \\ &\lesssim \sum_{E \in \mathcal{E}} h_E^{-1/2} \|\varphi_{\vec{\sigma}} - \mathcal{I}_{\text{grad}} \varphi_{\vec{\sigma}}\|_{L^2(E)} h_E^{1/2} \left\| \llbracket w_h \vec{n} \rrbracket \right\|_{H^1(E)} \lesssim \|\vec{\sigma}\|_{H(\text{div}, \Omega)} \left(\sum_{E \in \mathcal{E}} h_E \left\| \llbracket w_h \vec{n} \rrbracket \right\|_{H^1(E)}^2 \right)^{1/2}. \end{aligned}$$

Here, in the final line, we have also used the Cauchy–Schwarz inequality and (79b). The term t_2 can be estimated similarly:

$$\begin{aligned} t_2 &= \sum_{E \in \mathcal{E}} (\vec{\psi}_{\vec{\sigma}} - \mathcal{I}_{\text{div}} \vec{\psi}_{\vec{\sigma}}, \llbracket w_h \vec{n} \rrbracket)_E \leq \sum_{E \in \mathcal{E}} h_E^{-1/2} \|\vec{\psi}_{\vec{\sigma}} - \mathcal{I}_{\text{div}} \vec{\psi}_{\vec{\sigma}}\|_{L^2(E)} h_E^{1/2} \|\llbracket w_h \vec{n} \rrbracket\|_{L^2(E)} \\ &\lesssim \|\vec{\sigma}\|_{H(\text{div}, \Omega)} \left(\sum_{E \in \mathcal{E}} h_E \|\llbracket w_h \vec{n} \rrbracket\|_{L^2(E)}^2 \right)^{1/2}. \end{aligned}$$

Thus, the proof of the remaining inequality is complete:

$$\frac{\langle \vec{\sigma} \cdot \vec{n}, w_h \rangle_h^2}{\|\vec{\sigma}\|_{H(\text{div}, \Omega)}^2} = \frac{(t_1 + t_2)^2}{\|\vec{\sigma}\|_{H(\text{div}, \Omega)}^2} \lesssim \sum_{E \in \mathcal{E}} h_E \|\llbracket w_h \vec{n} \rrbracket\|_{L^2(E)}^2. \quad \square$$

Proof of Theorem 4.2. The DPG* solution (\vec{p}_h, v_h) satisfies the equations of (60) for all $(\vec{\tau}, \nu) \in \mathcal{V}_h$ and all $(\vec{\sigma}, \mu, \hat{\sigma}_n, \hat{\mu}) \in \mathcal{U}_h$. In particular, (60b) implies that $\langle \vec{p}_h \cdot \vec{n}, \hat{\mu} \rangle_h = \langle v_h, \hat{\sigma}_n \rangle_h = 0$. Hence the conditions of Lemmas 4.5 and 4.6 are satisfied. The conclusions of these lemmas show that the η in Theorem 4.1 satisfies

$$\eta((\vec{p}_h, v_h)) \approx \sum_{E \in \mathcal{E}_{\text{int}}} h_E \|\llbracket \vec{p}_h \cdot \vec{n} \rrbracket\|_{L^2(E)}^2 + \sum_{E \in \mathcal{E}} h_E \|\llbracket v_h \vec{n} \rrbracket\|_{H^1(E)}^2$$

and, moreover, the last term may be replaced by $h_E^{-1} \|\llbracket v_h \vec{n} \rrbracket\|_{L^2(E)}^2$, if we please. Hence an application of Theorem 4.1 completes the proof. \square

5. NUMERICAL EXPERIMENTS

In order to verify the mathematical theory developed above, we conducted several standard numerical verification experiments using two finite element software packages which have been used extensively for implementing DPG methods. In our first set of experiments, we used Camellia [46, 47], a user-friendly C++ toolbox developed by Nathan V. Roberts which itself relies on Sandia’s Trilinos library of packages [37]. Specifically, Camellia was used for the *a priori* convergence rate verification on the model square domain $\Omega_{\square} = [0, 1]^2$ reported on in Section 5.3. In our second set of experiments, we used *hp2D*, a sophisticated suite of Fortran routines with support for 2D local hierarchical and anisotropic *h*- and *p*-refinements on hybrid meshes [17] and corresponding oriented embedded shape functions for both quadrilateral and triangular elements in each of the canonical 2D de Rham sequence energy spaces [31]:

$$H^1(K) \xrightarrow{\text{grad}} H(\text{rot}, K) \xrightarrow{\text{rot}} L^2(K) \quad \text{and} \quad H^1(K) \xrightarrow{\text{curl}} H(\text{div}, K) \xrightarrow{\text{div}} L^2(K).$$

hp2D was used to implement a simple *hp*-adaptive algorithm for a singular solution to Poisson’s equation on the canonical L-shaped domain, $\Omega_{\mathbb{L}} = (-1, 1)^2 \setminus [0, 1] \times [-1, 0]$; see Section 5.4. This experiment parallels a similar study with the analogous DPG method in [19].

5.1. Set-up. Let $\Omega \in \{\Omega_{\square}, \Omega_{\mathbb{L}}\}$ and let Γ_D, Γ_N be disjoint and relatively open subsets comprising $\partial\Omega$; $\Gamma_D \cap \Gamma_N = \emptyset$, $\overline{\Gamma_D \cup \Gamma_N} = \partial\Omega$. All of our experiments investigate some form of Poisson’s equation:

$$(82) \quad \begin{cases} -\Delta v = f & \text{in } \Omega, \\ v = v_0 & \text{on } \Gamma_D, \\ \frac{\partial v}{\partial n} = p_n & \text{on } \Gamma_N, \end{cases}$$

where the load $f \in L^2(\Omega)$ and the boundary data v_0 and p_n are appropriately smooth.

As before, let Ω_h denote the mesh subordinate to Ω and let \mathcal{E} denote the corresponding collection of edges. In each of our experiments, we only considered piecewise-affine two-dimensional domains $\Omega \in \{\Omega_\square, \Omega_\boxplus\}$ subdivided into quadtree meshes consisting of either fully geometrically conforming or 1-irregular quadrilateral elements $K \in \Omega_h$.⁴ During stiffness matrix assembly, the degrees of freedom of every element edge with a hanging node was constrained by its common edge. Alternatively, because of the ultraweak variational formulation we considered, we could have incorporated each edge independently [46, 50].

For each quadrilateral element $K \in \Omega_h$, we associated a unique (anisotropic) polynomial order $p_K, q_K \geq 1$, respectively. Each associated polynomial order can be naturally related to a (p_K, q_K) -order conforming finite element de Rham sequence. For instance, begin with the standard Nédélec spaces of the first type,

$$\mathcal{Q}^{p_K, q_K}(K) \xrightarrow{\text{curl}} \mathcal{Q}^{p_K, q_K-1} \times \mathcal{Q}^{p_K-1, q_K}(K) \xrightarrow{\text{div}} \mathcal{Q}^{p_K-1, q_K-1}(K),$$

where $\mathcal{Q}^{p_K, q_K}(K)$ is the space of bivariate polynomials over K with degree at most p_K horizontally and q_K vertically. Now, consider the mesh-dependent sequence $W_{hp} \xrightarrow{\text{curl}} \mathbf{V}_{hp} \xrightarrow{\text{div}} Y_{hp}$, where

$$\begin{aligned} W_{hp} &= \{w \in H_0^1(\Omega) : w|_K \in \mathcal{Q}^{p_K, q_K}(K) \ \forall K \in \Omega_h\}, \\ \mathbf{V}_{hp} &= \{\vec{q} \in H(\text{div}, \Omega) : \vec{q}|_K \in \mathcal{Q}^{p_K, q_K-1}(K) \times \mathcal{Q}^{p_K-1, q_K}(K) \ \forall K \in \Omega_h\}, \\ Y_{hp} &= \{w \in L^2(\Omega) : w|_K \in \mathcal{Q}^{p_K-1, q_K-1}(K) \ \forall K \in \Omega_h\}. \end{aligned}$$

We define the (isotropic) uniform- p trial space to be $\mathcal{U}_h = Y_{hp} \times Y_{hp}^2 \times \text{tr}(W_{hp}) \times \text{tr}_n(\mathbf{V}_{hp})$, where $p_K = q_K = p$ is fixed for all $K \in \Omega_h$. Similarly, the corresponding (anisotropic) hp trial space is defined $\mathcal{U}_h = Y_{hp} \times Y_{hp}^2 \times \text{tr}(W_{hp}) \times \text{tr}_n(\mathbf{V}_{hp})$, where p_K and q_K are allowed to vary freely throughout the mesh. With the latter definition, notice that the polynomial order of an hp interface function, when restricted to a single shared edge $E \in \mathcal{E}$, $E = \bigcap_{\bar{K} \cap E \neq \emptyset} \bar{K}$, will naturally be restricted by the lowest polynomial order of all elements $\bar{K} \cap E \neq \emptyset$ sharing the edge.⁵

For the test functions, define the spaces

$$\begin{aligned} \widetilde{W}_{hp, dp} &= \{v \in H^1(\Omega_h) : v|_K \in \mathcal{Q}^{p_K+dp, q_K+dp}(K) \ \forall K \in \Omega_h\}, \\ \widetilde{\mathbf{V}}_{hp, dp} &= \{\vec{q} \in H(\text{div}, \Omega_h) : \vec{q}|_K \in \mathcal{Q}^{p_K+dp, q_K+dp-1}(K) \times \mathcal{Q}^{p_K+dp-1, q_K+dp}(K) \ \forall K \in \Omega_h\}. \end{aligned}$$

In all of our numerical experiments, we used $\mathcal{V}_h = \widetilde{W}_{hp, dp} \times \widetilde{\mathbf{V}}_{hp, dp}$ where $dp \in \{0, 1, 2\}$.

5.2. Adaptive mesh refinement. In our experiments with h - and hp -adaptive mesh refinement, we used a standard isotropic h -subdivision rule. Namely, at each refinement step, each element marked for h -refinement was uniformly subdivided into four equal-order quadrilateral elements. Afterward, a standard so-called “mesh closure” algorithm was called to induce a small number of additional isotropic h -subdivisions of neighboring elements in order to ensure 1-irregularity of the mesh. Alternatively, at each refinement step, the polynomial order of any p -refinement marked element was isotropically incremented by one, $(p_K, q_K) \mapsto (p_K+1, q_K+1)$,

⁴Although many of the preceding results are proven only for triangular meshes, the numerical experiments documented in this section verify alternative results in the setting of quadrilateral elements, which we understand to be similar.

⁵Such a mesh obeys the so-called *minimum rule*.

and then the order of all elements neighboring a p -refinement marked element was also isotropically incremented by one.

Recall that $\mathcal{E}_K = \{E \in \mathcal{E} : \text{meas}(\partial K \cap E) \neq \emptyset\}$ and define $\mathcal{E}_{K,\text{int}} = \mathcal{E}_K \cap \mathcal{E}_{\text{int}}$. Recalling the global error estimator $\eta_1(\vec{v}_h)$ appearing in [Theorem 4.2](#), define a *refinement indicator* $\eta_K \in \mathbb{R}_{\geq 0}$ for each $K \in \Omega_h$, viz.,

$$\eta_K = \left(\|\mathcal{L}\vec{v}_h - f\|_{L^2(K)}^2 + \sum_{E \in \mathcal{E}_{K,\text{int}}} h_E \|\llbracket \vec{p}_h \cdot \vec{n} \rrbracket\|_{L^2(E)}^2 + \sum_{E \in \mathcal{E}_K} h_E \|\llbracket v_h \vec{n} \rrbracket\|_{H^1(E)}^2 \right)^{1/2}.$$

In element marking, we followed the so-called “greedy” algorithm. That is, at each refinement step, all elements $K \in \Omega_h$ whose refinement indicator η_K was above 50% of the maximum over all elements in the mesh, $\eta_{\max} = \max_{K \in \Omega_h} \eta_K$, was marked for refinement. In the case of what we call h -adaptive mesh refinement, every marked element was h -refined, as described above, i.e. no elements were p -refined. Alternatively, in the case of hp -adaptive mesh refinement, a common flagging strategy [\[1\]](#) was used to decide whether to h or p refine; see [Section 5.4](#).

5.3. Pure Dirichlet boundary conditions on a square domain. Recall [\(82\)](#). In this first example, $\Omega = \Omega_{\square}$ and $\Gamma_D = \partial\Omega$. We considered two seemingly benign cases for the loads: (i) $f = 2\pi^2 \sin(\pi x) \sin(\pi y)$ and $v_0 = 0$; and (ii) $f = 0$ and $v_0 = 1$. In both cases, the exact solution is infinitely smooth. Indeed, in case (i), $v = \sin(\pi x) \sin(\pi y)$ and, in case (ii), $v = 1$. We first consider DPG* methods and then \mathcal{LL}^* methods.

5.3.1. The DPG* method. Recall from [Theorem 3.1](#) that the best approximation error of a DPG* method involves the Lagrange multiplier $\vec{\lambda} = (\vec{\zeta}, \lambda, \hat{\zeta}_n, \hat{\lambda})$ as well as the DPG* solution variable $\vec{v} = (\vec{p}, v)$. Assume that v is smooth. With the norm $\|(\vec{\tau}, \nu)\|_{\vec{V}}^2 = \|\vec{\tau}\|_{H(\text{div}, \Omega_h)}^2 + \|\nu\|_{H^1(\Omega_h)}^2$, λ solves


$$(83) \quad \begin{cases} -\Delta \lambda = g & \text{in } \Omega, \\ \lambda = 0 & \text{on } \partial\Omega, \end{cases}$$

where $g = v - 2\Delta v + \Delta^2 v$. Indeed, recall [\(62\)](#) and observe that $-\Delta \lambda = -\Delta f - \Delta e = \Delta(\Delta v) + (v + 2f) = v - 2\Delta v + \Delta^2 v$. Here, we have also used that $-\Delta v = f$ and $-\Delta e = v + 2f$. In case (i), $g = (1 + 4\pi^2 - 4\pi^4) \sin(\pi x) \sin(\pi y)$, meanwhile, in case (ii), $g = 1$. Notice that $g \in C^\infty(\overline{\Omega})$ in both cases.



In the first case, λ can easily be shown to be a constant scalar multiple of $\sin(\pi x) \sin(\pi y)$ and so $\lambda \in C^\infty(\overline{\Omega})$ is infinitely smooth. Therefore, by [Corollary 3.4](#), the convergence rate of the DPG* method under uniform h -refinement will be limited only by the underlying de Rham sequence polynomial order p . Indeed, [Figure 5.5 \(A\)](#) demonstrates the convergence of the corresponding discrete solution $\vec{v}_h = (\vec{p}_h, v_h)$ to the exact solution, $\vec{v} = (\text{grad } v, v)$, measured in the full test norm above, starting with an single-element mesh with (isotropic) polynomial order $p_K = q_K = p \in \{1, 2, 3, 4\}$. [Figure 5.5 \(B\)](#) presents the convergence of only the solution variable v_h , measured in the $L^2(\Omega)$ -norm. Although both figures correspond only to a test space enrichment of $dp = 1$, similar results were observed for each choice $dp \in \{0, 1, 2\}$.

In the second case, due to the shape of the domain, we can guarantee that $\lambda \in H^3(\Omega)$ but, ultimately, the corresponding Lagrange multiplier $\lambda \notin C^\infty(\overline{\Omega})$ is still *not* infinitely smooth.⁶ Therefore, with this problem, the DPG* method experiences rate-limited convergence


⁶Standard elliptic regularity theory can be used to show that $\lambda \in C^\infty(\Omega)$ is, however, infinitely smooth in the interior of the domain [\[29\]](#).



(A) Optimal rates.

(B) Optimal rates in the L^2 -norm.FIGURE 5.1. Convergence under h -uniform mesh refinements with the manufactured solution $v(x, y) = \sin(\pi x) \sin(\pi y)$. (Here, $dp = 1$.)

(A) Limited optimal rates.

(B) Limited optimal rates in the L^2 -norm.FIGURE 5.2. Convergence under h -uniform mesh refinements with the manufactured solution $v(x, y) = 1$. (Here, $dp = 1$.)

under uniform h -refinements. This is evidenced by Figure 5.2. However, as demonstrated by Figure 5.3, using the greedy h -refinement strategy from Section 5.2, optimal convergence rates can still be recovered through adaptive mesh refinement. See Figure 5.4 for a visual depiction of the solution of the corresponding auxiliary problem (83) as well as the corresponding adaptively refined mesh.

Remark 5.1. Previously in this subsection, we remarked that similar results were observed for each test space enrichment parameter $dp \in \{0, 1, 2\}$ that we chose in our numerical experiments analyzing case (i). Similarly, in case (ii), the behaviour documented above was nearly indistinguishable for each $dp \geq 1$. However, when $dp = 0$ we observed unexpected effects which we repeatedly verified with independent implementations of the method. Indeed, starting from a mesh consisting of a single square element of order p and subsequently performing uniform h -refinements, the exact solution $v = 1$ was repeatedly reproduced up to machine zero, no matter the polynomial order $p \in \{1, 2, 3, 4\}$ considered. In testing more complicated manufactured solutions (not documented here) which also feature a singular Lagrange multiplier λ , we discovered superconvergence effects from this choice of enrichment parameter. Indeed, in our numerous additional verification experiments with dp set to zero, the method overcame the




FIGURE 5.3. Convergence under h -adaptive mesh refinements with the manufactured solution $v(x, y) = 1$. (Here, $dp = 1$.)




FIGURE 5.4. Left: The DPG* solution variable λ when $v = 1$. (Color scale represents the solution values.) Right: The corresponding quadtree mesh coming from the h -adaptive algorithm after ten refinements. (Here, $p = 4$ and $dp = 1$.)

rate-limited behavior illustrated in Figure 5.2. This peculiar superconvergence artifact can not be explained by the theory presented in this paper. Notably, this artifact also agrees with previous results seen with a DPG* method for acoustics which can be found in the original technical report on the method [40] (which portions of this text are based off of) and clearly warrants further analysis.

5.3.2. Comparison with the \mathcal{LL}^* method. Consider the operator $\mathcal{L}: H(\text{div}, \Omega) \times H^1(\Omega) \rightarrow (L^2(\Omega))^d \times L^2(\Omega)$ defined in (41), and its adjoint, $\mathcal{L}^*: H(\text{div}, \Omega) \times H^1(\Omega) \rightarrow (L^2(\Omega))^d \times L^2(\Omega)$, defined in (42):

$$\mathcal{L}(\vec{p}, v) = (\vec{p} - \text{grad } v, -\text{div } \vec{p}), \quad \mathcal{L}^*(\vec{\sigma}, \mu) = (\vec{\sigma} + \text{grad } \mu, \text{div } \vec{\sigma}).$$

One may write an \mathcal{LL}^* formulation of Poisson's equation (82), with $\Gamma_D = \partial\Omega$, in the following way:

$$(84) \quad (\mathcal{L}^*(\vec{\zeta}, \lambda), \mathcal{L}^*(\vec{\sigma}, \mu))_\Omega = (f, \mu)_\Omega + \langle v_0, \vec{\sigma} \cdot \vec{n} \rangle_{H^{-1/2}(\partial\Omega)},$$

for all $(\vec{\sigma}, \mu) \in H(\text{div}, \Omega) \times H_0^1(\Omega)$. After solving (84) for the unique $(\vec{\zeta}, \lambda) \in H(\text{div}, \Omega) \times H_0^1(\Omega)$, the solution of (82) may be recovered by simply applying the adjoint operator to the Lagrange multiplier; $\mathcal{L}^*(\vec{\zeta}, \lambda) = (\vec{p}, v) = (\text{grad } v, v)$.

It is instructive to write out the strong form of the equations for $\vec{\zeta}$ and λ , which are implied by (84):


$$(85) \quad \begin{cases} \vec{\zeta} + \text{grad } \lambda - \text{grad } \text{div } \vec{\zeta} = \vec{0} & \text{in } \Omega, \\ -\text{div } \vec{\zeta} - \Delta \lambda = f & \text{in } \Omega, \\ \lambda = 0 & \text{on } \partial\Omega, \\ \text{div } \vec{\zeta} = v_0 & \text{on } \partial\Omega. \end{cases}$$

By simply substituting the following expressions into (85), the reader may verify that the unique Lagrange multipliers $\vec{\zeta}$ and λ can also be written


$$-\Delta \lambda = f + v, \quad \vec{\zeta} = -\text{grad } \lambda + \text{grad } v,$$

where v is just the unique solution of (82).

We can now see that the \mathcal{LL}^* method has precisely the same rate-limited behavior as the DPG* method. Indeed, consider the following two loading scenarios again: (i) $f = 2\pi^2 \sin(\pi x) \sin(\pi y)$ and $v_0 = 0$; and (ii) $f = 0$ and $v_0 = 1$. In case (i), both $v, \lambda \in C^\infty(\bar{\Omega})$ are infinitely smooth. On the other hand, in case (ii), $\lambda \in H^3(\Omega)$ but $\lambda \notin C^\infty(\bar{\Omega})$, even though the solution of (82), $v = 1$, is just a constant function. The numerical evidence in Figure 5.5 demonstrates that the convergence rate of the \mathcal{LL}^* method is affected in case (ii). One may also compare this outcome with [10, Theorem 6.1], which guarantees optimal convergence rates *only* under the assumption that the Lagrange multiplier is sufficiently smooth.



(A) Optimal rates; $v(x, y) = \sin(\pi x) \sin(\pi y)$.



(B) Limited optimal rates; $v(x, y) = 1$.

FIGURE 5.5. Convergence under h -uniform mesh refinements with the \mathcal{LL}^* method.

Remark 5.2. Many \mathcal{LL}^* formulations contain additional terms involving a (two- or three-dimensional) curl operator. Such terms arise due to the general fact that $\text{curl } \vec{p} = \text{curl } \text{grad } v = \vec{0}$, thus allowing $H(\text{div}, \Omega)$ to be replaced by $H(\text{div}, \Omega) \cap H(\text{curl}, \Omega) \subseteq (H^1(\Omega))^d$ in the domains of the differential operators above. This space facilitates the use of C^0 piecewise polynomial finite element bases when the solution is sufficiently regular. Nevertheless, enforcing the additional equation $\text{curl } \vec{p} = \vec{0}$ also requires introducing additional “slack” variables which make the ultimate expression for the new operators \mathcal{L} and \mathcal{L}^* much more complicated. The analysis above can easily be repeated in such scenarios, but the outcome of limited convergence rates will not change.

5.4. Mixed boundary conditions on an L-shaped domain. Again, recall (82). In this final example, set $\Omega = \Omega_{\mathbb{P}}$, $\Gamma_D = [0, 1] \times \{0\} \cup \{0\} \times [0, -1]$, and $v_0 = 0$. Additionally, set p_n to be the normal derivative of the exact solution $v(r, \theta) = r^{2/3} \sin(\frac{2}{3}\theta)$. For this problem, it is well known that the solution $v \in H^{1+s}(\Omega)$, for all $s < 2/3$.

In each of our experiments, we began with a single three-element mesh composed of congruent squares and uniform order $p_K = q_K = 2$ and $dp = 1$ in all three elements. Figure 5.6 (A) demonstrates the convergence of the solution error we witnessed under h -uniform, h -adaptive (as described above), and hp -adaptive refinements using a flagging strategy where all marked element adjacent to the origin (i.e. the singular point) are h -refined and all other marked elements are p -refined. As shown in Figure 5.6 (B), the error estimator $\eta(\vec{v}_h)$ generally overestimated the solution error and the dependence upon dp was not seen to be roundly significant. Figure 5.7 depicts both the computed solution and the hp mesh mesh after fifteen refinement steps.




FIGURE 5.6. (A): Comparison of the convergence of various refinement strategies when $dp = 1$. (B): Convergence of the error in the DPG* solution variable \vec{v}_h and the error estimator $\eta(\vec{v}_h)$ for two values of dp with the hp -adaptive algorithm.




FIGURE 5.7. Left: The DPG* solution v . (Color scale represents solution values.) Right: The corresponding hp quadtree mesh found by the hp -adaptive algorithm after fifteen refinements. (Colors represent polynomial degrees p .)

REFERENCES

- [1] M. Ainsworth and B. Senior. Aspects of an adaptive hp -finite element method: Adaptive strategy, conforming approximation and efficient solvers. *Comput. Methods Appl. Mech. Engrg.*, 150(1-4):65–87, 1997.
- [2] C. Bacuta and K. Qirko. A saddle point least squares approach to mixed methods. *Comput. Math. Appl.*, 70(12):2920 – 2932, 2015.
- [3] Å. Björck. *Numerical methods for least squares problems*. SIAM, 1996.
- [4] P. B. Bochev and M. D. Gunzburger. *Least-squares finite element methods*, volume 166. Springer Science & Business Media, 2009.
- [5] D. Boffi, M. Fortin, and F. Brezzi. *Mixed finite element methods and applications*. Springer series in computational mathematics. Springer, Berlin, Heidelberg, 2013.
- [6] T. Bouma, J. Gopalakrishnan, and A. Harb. Convergence rates of the DPG method with reduced test space degree. *Comput. Math. Appl.*, 68:1550–1561, 2014.
- [7] J. Bramwell, L. Demkowicz, J. Gopalakrishnan, and Q. Weifeng. A locking-free hp DPG method for linear elasticity with symmetric stresses. *Numer. Math.*, 122(4):671–707, 2012.
- [8] D. Broersen and R. P. Stevenson. A Petrov–Galerkin discretization with optimal test space of a mild-weak formulation of convection–diffusion equations in mixed form. *IMA J. Numer. Anal.*, 35(1):39–73, 2015.
- [9] T. Bui-Thanh, L. Demkowicz, and O. Ghattas. A unified discontinuous Petrov–Galerkin method and its analysis for Friedrichs’ systems. *SIAM J. Numer. Anal.*, 51(4):1933–1958, 2013.
- [10] Z. Cai, T. A. Manteuffel, S. F. McCormick, and J. Ruge. First-order system \mathcal{LL}^* (FOSLL*): Scalar elliptic partial differential equations. *SIAM J. Numer. Anal.*, 39(4):1418–1445, 2001.
- [11] C. Carstensen, L. Demkowicz, and J. Gopalakrishnan. Breaking spaces and forms for the DPG method and applications including Maxwell equations. *Comput. Math. Appl.*, 72(3):494–522, 2016.
- [12] C. Carstensen and F. Hellwig. Low-order discontinuous Petrov–Galerkin finite element methods for linear elasticity. *SIAM J. Numer. Anal.*, 54(6):3388–3410, 2016.
- [13] C. Carstensen and S. Puttkammer. A low-order discontinuous Petrov–Galerkin method for the Stokes equations. *Numer. Math.*, 140(1):1–34, 2018.
- [14] J. Chan, N. Heuer, T. Bui-Thanh, and L. Demkowicz. A robust DPG method for convection-dominated diffusion problems II: Adjoint boundary conditions and mesh-dependent test norms. *Comput. Math. Appl.*, 67(4):771–795, 2014.
- [15] P. Clément. Approximation by finite element functions using local regularization. *Rev. fr. autom. inform. rech. opér., Anal. numér.*, 9(R2):77–84, 1975.
- [16] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and variational stabilization for convection-diffusion equations. *ESAIM Math. Model. Numer. Anal.*, 46(5):1247–1273, 2012.
- [17] L. Demkowicz. *Computing with hp Finite Elements. I. One and Two Dimensional Elliptic and Maxwell Problems*. Chapman & Hall/CRC Press, New York, October 2006.
- [18] L. Demkowicz and J. Gopalakrishnan. A class of discontinuous Petrov–Galerkin methods. Part I: The transport equation. *Comput. Methods Appl. Mech. Engrg.*, 199(23-24):1558–1572, 2010.
- [19] L. Demkowicz and J. Gopalakrishnan. Analysis of the DPG method for the Poisson equation. *SIAM J. Numer. Anal.*, 49(5):1788–1809, 2011.
- [20] L. Demkowicz and J. Gopalakrishnan. A class of discontinuous Petrov–Galerkin methods. II. Optimal test functions. *Numer. Methods Partial Differ. Equ.*, 27(1):70–105, 2011.
- [21] L. Demkowicz and J. Gopalakrishnan. A primal DPG method without a first-order reformulation. *Comput. Math. Appl.*, 66(6):1058–1064, 2013.
- [22] L. Demkowicz and J. Gopalakrishnan. Discontinuous Petrov–Galerkin (DPG) method. In E. Stein, R. Borst, and T. J. R. Hughes, editors, *Encyclopedia of Computational Mechanics Second Edition*, pages 1–15. Wiley Online Library, 2017.
- [23] L. Demkowicz, J. Gopalakrishnan, S. Nagaraj, and P. Sepulveda. A spacetime DPG method for the Schrödinger equation. *SIAM J. Numer. Anal.*, 55(4):1740–1759, 2017.
- [24] L. Demkowicz, J. Gopalakrishnan, and J. Schöberl. Polynomial extension operators. Part III. *Math. Comput.*, 81(279):1289–1326, 2012.
- [25] L. Demkowicz and N. Heuer. Robust DPG method for convection-dominated diffusion problems. *SIAM J. Numer. Anal.*, 51(5):2514–2537, 2013.

- [26] A. Demlow and A. N. Hirani. A posteriori error estimates for finite element exterior calculus: The de Rham complex. *Found. Computat. Math.*, 14(6):1337–1371, 2014.
- [27] A. Ern, J.-L. Guermond, and G. Caplain. An intrinsic criterion for the bijectivity of Hilbert operators related to Friedrichs’ systems. *Comm. Partial Differential Equations*, 32(1-3):317–341, 2007.
- [28] J. Ernesti. *Space-Time Methods for Acoustic Waves with Applications to Full Waveform Inversion*. PhD thesis, Karlsruhe, Germany, 2017.
- [29] L. C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [30] F. Fuentes, B. Keith, L. Demkowicz, and P. Le Tallec. Coupled variational formulations of linear elasticity and the DPG methodology. *J. Comput. Phys.*, 348:715–731, 2017.
- [31] F. Fuentes, B. Keith, L. Demkowicz, and S. Nagaraj. Orientation embedded high order shape functions for the exact sequence elements of all shapes. *Comput. Math. Appl.*, 70(4):353–458, 2015.
- [32] T. Führer. Superconvergence in a DPG method for an ultra-weak formulation. *Comput. Math. Appl.*, 75(5):1705–1718, 2018.
- [33] T. Führer. Superconvergent DPG methods for second-order elliptic problems. *Comput. Meth. Appl. Mat.*, 19(3):483–502, 2019.
- [34] J. Gopalakrishnan, I. Muga, and N. Olivares. Dispersive and dissipative errors in the DPG method with scaled norms for Helmholtz equation. *SIAM J. Sci. Comput.*, 36(1):A20–A39, 2014.
- [35] J. Gopalakrishnan and W. Qiu. An analysis of the practical DPG method. *Math. Comput.*, 83(286):537–552, 2014.
- [36] J. Gopalakrishnan and P. Sepulveda. A spacetime DPG method for acoustic waves. In U. Langer and O. Steinbach, editors, *Space-Time Methods: Applications to Partial Differential Equations*, pages 129–154. De Gruyter, Berlin, Boston, 2019.
- [37] M. A. Heroux, R. A. Bartlett, V. E. Howle, R. J. Hoekstra, J. J. Hu, T. G. Kolda, R. B. Lehoucq, K. R. Long, R. P. Pawlowski, E. T. Phipps, A. G. Salinger, H. K. Thornquist, R. S. Tuminaro, J. M. Willenbring, A. Williams, and K. S. Stanley. An overview of the Trilinos project. *ACM Trans. Math. Softw.*, 31(3):397–423, 2005.
- [38] B. Keith. *New ideas in adjoint methods for PDEs: A saddle-point paradigm for finite element analysis and its role in the DPG methodology*. PhD thesis, The University of Texas at Austin, Austin, Texas, U.S.A., 2018.
- [39] B. Keith, A. V. Astaneh, and L. Demkowicz. Goal-oriented adaptive mesh refinement for discontinuous Petrov–Galerkin methods. *SIAM J. Numer. Anal.*, 57(4):1649–1676, 2019.
- [40] B. Keith, L. Demkowicz, and J. Gopalakrishnan. DPG* method. ICES Report 17-25, The University of Texas at Austin, 2017.
- [41] B. Keith, F. Fuentes, and L. Demkowicz. The DPG methodology applied to different variational formulations of linear elasticity. *Comput. Methods Appl. Mech. Engrg.*, 309:579–609, 2016.
- [42] B. Keith, S. Petrides, F. Fuentes, and L. Demkowicz. Discrete least-squares finite element methods. *Comput. Methods Appl. Mech. Engrg.*, 327:226–255, 2017.
- [43] S. Nagaraj, S. Petrides, and L. F. Demkowicz. Construction of DPG Fortin operators for second order problems. *Comput. Math. Appl.*, 74(8):1964–1980, 2017.
- [44] S. Petrides and L. F. Demkowicz. An adaptive DPG method for high frequency time-harmonic wave propagation problems. *Comput. Math. Appl.*, 74(8):1999–2017, 2017.
- [45] S. Repin, S. Sauter, and A. Smolianski. Two-sided a posteriori error estimates for mixed formulations of elliptic problems. *SIAM J. Numer. Anal.*, 45(3):928–945, 2007.
- [46] N. V. Roberts. Camellia: A software framework for discontinuous Petrov–Galerkin methods. *Comput. Math. Appl.*, 68(11):1581–1604, 2014.
- [47] N. V. Roberts. Camellia v1.0 manual: Part I. Technical Report ANL/ALCF-16/3, Argonne National Laboratory, Argonne, Illinois, 2016.
- [48] N. V. Roberts, T. Bui-Thanh, and L. Demkowicz. The DPG method for the Stokes problem. *Comput. Math. Appl.*, 67(4):966–995, 2014.
- [49] J. Schöberl. A posteriori error estimates for Maxwell equations. *Math. Comput.*, 77(262):633–649, 2008.
- [50] A. Vaziri Astaneh, F. Fuentes, J. Mora, and L. Demkowicz. High-order polygonal discontinuous Petrov–Galerkin (PolyDPG) methods using ultraweak formulations. *Comput. Methods Appl. Mech. Engrg.*, 332:686–711, 2017.

- [51] A. Vaziri Astaneh, B. Keith, and L. Demkowicz. On perfectly matched layers for discontinuous Petrov–Galerkin methods. *Comput. Mech.*, 63(6):1131–1145, 2019.
- [52] R. Verfürth. *A review of a posteriori error estimation and adaptive mesh-refinement techniques*. John Wiley & Sons Inc, 1996.
- [53] C. Wieners. The skeleton reduction for finite element substructuring methods. In *Numerical Mathematics and Advanced Applications ENUMATH 2015*, pages 133–141. Springer, 2016.

(Demkowicz) THE INSTITUTE FOR COMPUTATIONAL ENGINEERING AND SCIENCES (ICES),
THE UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TX 78712, U.S.A.

Email address: `leszek@ices.utexas.edu`

(Gopalakrishnan) PORTLAND STATE UNIVERSITY, PORTLAND, OR 97207-0751, U.S.A.

Email address: `gjay@pdx.edu`

(Keith) TECHNISCHE UNIVERSITÄT MÜNCHEN, GARCHING, 85748, GERMANY

Email address: `keith@ma.tum.de`