



Dispersive Ordering Is the Same as Tail-Ordering

Author(s): J. V. Deshpande and S. C. Kocher

Source: *Advances in Applied Probability*, Vol. 15, No. 3 (Sep., 1983), pp. 686-687

Published by: [Applied Probability Trust](#)

Stable URL: <http://www.jstor.org/stable/1426626>

Accessed: 13/11/2010 22:50

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=apt>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Applied Probability Trust is collaborating with JSTOR to digitize, preserve and extend access to *Advances in Applied Probability*.

<http://www.jstor.org>

LETTERS TO THE EDITOR

DISPERSIVE ORDERING IS THE SAME AS TAIL-ORDERING

J. V. DESHPANDE* AND
 S. C. KOCHAR,* *Panjab University*

Recently Lewis and Thompson (1981) and Shaked (1982) have defined the following partial ordering on the space of probability distribution functions. Let F and G be two cumulative distribution functions with F^{-1} and G^{-1} as the respective left-continuous inverses. Then F is said to be dispersed with respect to G ($F \stackrel{\text{disp}}{<} G$) if and only if

$$(1) \quad F^{-1}(\beta) - F^{-1}(\alpha) \leq G^{-1}(\beta) - G^{-1}(\alpha) \quad \text{whenever } 0 < \alpha < \beta < 1.$$

In these papers many interesting and important properties of this ordering have been studied. We wish to point out some connections that this ordering has with some concepts extensively used in reliability theory and related statistical inference.

It is easily seen by putting $\alpha = F(x)$ and $\beta = F(y)$, where $x \leq y$ that $F \stackrel{\text{disp}}{<} G$ if and only if

$$(2) \quad G^{-1}F(x) - x \text{ is non-decreasing in } x.$$

In the literature (see Doksum (1969)) Condition (2) has already been used. F is said to be tail-ordered with respect to G ($F \stackrel{t}{<} G$) if and only if (2) holds. Doksum (1969) has also studied some properties of this ordering. Yanagimoto and Sibuya (1976), (1980) call this partial ordering (introduced originally by Fraser (1957)), ' G is stochastically more spread than F ,' and have discussed some properties of this ordering. *Thus we see that dispersive ordering is the same as tail-ordering.* Doksum (1969), Deshpande and Kochar (1982) and Deshpande and Mehta (1982) have used this concept of tail-ordering in some inferential problems to obtain bounds on efficiencies of tests, probabilities of correct selections, etc.

For distributions with $F(0) = G(0) = 0$ another partial ordering is given in Shaked (1982). It is characterized by

$$(3) \quad \frac{G^{-1}(\beta)}{G^{-1}(\alpha)} \geq \frac{F^{-1}(\beta)}{F^{-1}(\alpha)}, \quad \text{whenever } 0 < \alpha < \beta < 1.$$

Again substituting $x = F^{-1}(\alpha)$ and $y = F^{-1}(\beta)$, $x \leq y$, we see that (3) is true if and only if

$$(4) \quad \frac{G^{-1}F(x)}{x} \text{ is non-decreasing in } x,$$

as noted by Yanagimoto and Sibuya as early as 1976.

Received 7 March 1983; revision received 19 May 1983.

* Postal address: Department of Statistics, Panjab University, Chandigarh, India.

This is used in the literature to define the well-known concept of star-ordering. F is said to be star-ordered with respect to $G (F \stackrel{*}{\leq} G)$ if $G^{-1} F(x)$ is a star-shaped function on $(0, \infty)$ or in other words $G^{-1}F(x)/x$ is non-decreasing (see e.g. Barlow and Proschan (1975)). Thus the results of Saunders and Moran (1978) and Shaked (1982) regarding the gamma family parametrized by the shape parameter being ordered according to (3) are the same as the well-known result that it is ordered according to star-ordering. In fact, it is known to be ordered according to the stronger convex-ordering ($G^{-1}F(x)$ is a convex function of x) as well.

Some of the other results proved in Lewis and Thompson (1981) and Shaked (1982), when stated in terms of tail-ordering and star-ordering are perhaps known to the workers in the area. A connected result first proved in Kochar (1978) is given below.

Theorem. Let F and G be absolutely continuous such that $F(0) = G(0) = 0$ and let the corresponding density functions be such that $f(0) \geq g(0) > 0$. Then $f \stackrel{*}{\leq} G$ implies $F \stackrel{t}{\leq} G$.

Proof.

$$(5) \quad F \stackrel{*}{\leq} G \Leftrightarrow \frac{G^{-1}F(x)}{x} \text{ is non-decreasing in } x$$

$$\Leftrightarrow \frac{f(x)}{g[G^{-1}F(x)]} \geq \frac{G^{-1}F(x)}{x} \text{ for } x \geq 0$$

by differentiation. Now since

$$\lim_{x \rightarrow 0^+} \frac{G^{-1}F(x)}{x} = \frac{f(0)}{g(0)} \geq 1,$$

and $G^{-1}F(x)/x$ is non-decreasing, (5) implies $f(x) \geq g[G^{-1}F(x)], x > 0$

$$\Leftrightarrow F \stackrel{t}{\leq} G.$$

The last step is observed also in Shaked (1982), Remark 2.3.

References

- BARLOW, R. E. AND PROSCHAN, F. (1975) *Statistical Theory of Reliability and Life Testing*. Holt, Rinehart and Winston, New York.
- DESHAPANDE, JAYANT V. AND KOCHAR, SUBHASH C. (1982) Some competitors of Wilcoxon-Mann-Whitney test for the location alternative. *J. Indian Statist. Assoc.* **19**, 9–18.
- DESHAPANDE, JAYANT V. AND MEHTA, GOBIND P. (1982) Inequality for the infimum of PCS for heavy tailed distributions. *J. Indian Statist. Assoc.* **19**, 19–25.
- DOKSUM, K. (1969) Star-shaped transformations and the power of rank tests. *Ann. Math. Statist.* **40**, 1167–1176.
- FRASER, D. A. S. (1957) *Nonparametric Methods in Statistics*. Wiley, New York.
- KOCHAR, SUBHASH C. (1978) Distribution-free Comparison of Two Probability Distributions with Reference to their Hazard Rates. Unpublished Ph.D Thesis, Panjab University, India.
- LEWIS, T. AND THOMPSON, J. W. (1981) Dispersive distributions and the connection between dispersivity and strong unimodality. *J. Appl. Prob.* **18**, 76–90.
- SAUNDERS, I. W. AND MORAN, P. A. P. (1978) On the quantiles of the gamma and F distributions. *J. Appl. Prob.* **15**, 426–432.
- SHAKED, M. (1982) Dispersive ordering of distributions. *J. Appl. Prob.* **19**, 310–320.
- YANAGIMOTO, T. AND SIBUYA, M. (1976) Isotonic tests for spread and tail. *Ann. Inst. Statist. Math.* **28** A, 329–342.
- YANAGIMOTO, T. AND SIBUYA, M. (1980) Comparison of tails of distributions in models for estimating safe doses. *Ann. Inst. Statist. Math.* **32** A, 325–340.