

SOME BIVARIATE NOTIONS OF IFR AND DMRL AND RELATED PROPERTIES

BRUNO BASSAN,* ** *Università di Roma 'La Sapienza'*

SUBHASH KOCHAR,*** *Indian Statistical Institute*

FABIO SPIZZICHINO,* *Università di Roma 'La Sapienza'*

Abstract

Recently Bassan and Spizzichino (1999) have given some new concepts of multivariate ageing for exchangeable random variables, such as a special type of bivariate IFR, by comparing distributions of residual lifetimes of dependent components of different ages. In the same vein, we further study some properties of these concepts of IFR in the bivariate case. Then we introduce certain concepts of bivariate DMRL ageing and we develop a treatment that parallels those developed for bivariate IFR. For both the IFR and DMRL concepts, we analyse a weak and a strong version, and discuss some of the differences between them.

Keywords: Residual lifetime; mean residual life function; multivariate extensions of IFR and DMRL; log-concavity; Schur-concavity; ageing properties of mixtures

AMS 2000 Subject Classification: Primary 62N05; 60G09; 90B25

1. Introduction

Univariate concepts of ageing such as IFR (increasing failure rate), NBU (new better than used), DMRL (decreasing mean residual life) have played an important role in survival analysis, reliability theory, maintenance policies, operations research and many other areas of applied probability. They have also been found useful in getting bounds and inequalities on efficiencies of estimates and tests.

A complex system usually consists of several components which are working under the same environment and hence their lifetimes are, generally, dependent. In the literature, several attempts have been made to extend the concepts of univariate ageing to the multivariate case. Some important references are Brindley and Thompson (1972), Arjas (1981), Savits (1985), Shaked and Shanthikumar (1988), (1991), Barlow and Mendel (1993), Barlow and Spizzichino (1993) and Bassan and Spizzichino (1999), among others. In this paper, we consider the case when the lifetimes of the components have exchangeable joint probability distribution. We introduce some new notions of multivariate ageing and also discuss further the properties of those other notions recently introduced in the literature. We fix our attention only on the bivariate case, though these ideas can be easily extended to the multivariate case. We shall be assuming that the random variables under consideration are nonnegative. Absolute continuity will be tacitly assumed whenever needed.

Received 3 January 2002; revision received 15 May 2002.

* Postal address: Dipartimento di Matematica, Università di Roma 'La Sapienza', Piazzale Aldo Moro 5, 00185 Roma, Italy.

** Email address: bassan@mat.uniroma1.it

*** Postal address: Indian Statistical Institute, 7 S. J. S. Sansanwal Marg, New Delhi 110016, India.

Both from the technical and conceptual point of view, the interest in the paper is concentrated on the conditional distributions of residual lifetimes, given different types of survival data. In fact, these objects will play a central role both in the formulation of the different definitions and in the interpretation of related results.

Before we go into the details, let us quickly review some common notions of univariate positive ageing and stochastic orderings of various kinds. We shall denote the density function, the survival function and the hazard rate function of a univariate random variable X by f_X , \bar{F}_X and r_{F_X} respectively.

A random variable X is said to be *stochastically* larger than another random variable Y (denoted by $X \geq_{st} Y$) if $\bar{F}_X(x) \geq \bar{F}_Y(x)$ for all x . A stronger notion of stochastic dominance is that of *hazard rate* ordering. A random variable X is said to be larger than Y in *hazard rate* ordering (denoted by $X \geq_{hr} Y$) if $\bar{F}_X(x)/\bar{F}_Y(x)$ is nondecreasing in x . In the continuous case, this is equivalent to $r_{F_X}(x) \leq r_{F_Y}(x)$ for all x . Finally, X is said to be larger than Y in *likelihood ratio* ordering (denoted by $X \geq_{lr} Y$) if $f_X(x)/f_Y(x)$ is nondecreasing in x . In case X and Y have a common left end-point of their supports, we have the following chain of implications among the above stochastic orders:

$$X \geq_{lr} Y \implies X \geq_{hr} Y \implies X \geq_{st} Y.$$

Let X denote the lifetime of a unit with survival function \bar{F} and let us denote by \bar{F}_t the survival function of the residual life of a unit surviving at time t , that is,

$$\begin{aligned} \bar{F}_t(x) &= P[X - t > x \mid X > t] \\ &= \frac{\bar{F}(x+t)}{\bar{F}(t)}. \end{aligned}$$

Let us denote by X_t a random variable with survival function \bar{F}_t . A unit with survival function \bar{F} ages positively with time if the random variable X_t decreases with time t in some stochastic sense. That is, if

$$X_{t_1} \geq_* X_{t_2},$$

for all (t_1, t_2) in some region $A \subset \{(t_1, t_2) \mid 0 \leq t_1 < t_2\}$ where \geq_* can be any of the above one-dimensional stochastic orderings \geq_{st} , \geq_{hr} , \geq_{lr} or the orderings of the means. Different concepts of positive ageing correspond to different regions A and different stochastic orders between X_{t_1} and X_{t_2} for $(t_1, t_2) \in A$ (see the discussion in Spizzichino (2001)). For example, a random variable X is NBU if and only if $X \geq_{st} X_t$ for $t > 0$. Furthermore, a random variable X has IFR distribution if and only if $X_{t_1} \geq_{st} X_{t_2}$ for $t_1 < t_2$ and also if and only if $X_{t_1} \geq_{hr} X_{t_2}$ for $t_1 < t_2$.

The mean residual life (MRL) of a unit or a subject at age t is the average remaining life among those population members who have survived until time t . The mean residual life function (MRLF) is defined as

$$\begin{aligned} \mu_F(t) &= E[X_t] \\ &= E[X - t \mid X > t] \\ &= \int_0^\infty \bar{F}_t(x) dx \\ &= \frac{\int_t^\infty \bar{F}(x) dx}{\bar{F}(t)}. \end{aligned}$$

Like the failure rate function, the MRLF can be used to describe conditional concepts of ageing; however, the MRLF is more intuitive, especially in the health sciences. The review article by Guess and Proschan (1988) gives a nice summary of the theory of MRL and an extensive bibliography.

In many applications it is reasonable to assume that the life system is monotonically degenerating. This concept has been modelled several ways, of which increasing failure rate (IFR) is probably the most studied. The somewhat weaker version of decreasing mean residual life (DMRL), which is implied by IFR, is perhaps more clear conceptually and is easier to explain to the user.

A random variable X is said to have a DMRL (decreasing mean residual life) distribution if $\mu_F(t) = E[X_t]$ is decreasing in t . Note that X is DMRL if and only if the function $\int_t^\infty \bar{F}(x) dx$ is log-concave in t . Kochar *et al.* (2000) consider the problem of estimating the MRLF of a DMRL distribution.

A weaker concept of positive ageing is that of new better than used in expectation (NBUE). A random variable X is said to have NBUE distribution if

$$\mu_F(t) \leq \mu_F(0) \quad \text{for } t > 0.$$

See Barlow and Proschan (1981) and Deshpande *et al.* (1986) for the definitions of other concepts of ageing.

Let $\mu_{F_X}(t)$ and $\mu_{F_Y}(t)$ denote the mean residual life functions of X and Y respectively.

Definition 1.1. The random variable X is said to be greater than Y according to mean residual life (MRL) ordering (written as $X \geq_{\text{mrl}} Y$) if

$$\mu_{F_X}(t) \geq \mu_{F_Y}(t) \quad \text{for } t \geq 0.$$

It is immediately obvious that $X \geq_{\text{mrl}} Y$ if and only if

$$\frac{\int_t^\infty \bar{F}(x) dx}{\int_t^\infty \bar{G}(x) dx}$$

is nondecreasing in t .

Note that hazard rate ordering implies mean residual life ordering, but the converse is not true. For more details on stochastic orderings, see Chapter 1 of Shaked and Shanthikumar (1994).

The organization of the paper is as follows. In Section 2, we recall the concept of bivariate IFR ageing as introduced by Bassan and Spizzichino (1999) and study some of its new properties. In Section 3, we introduce the concept of bivariate DMRL distributions for exchangeable random variables and study its properties. In Appendix A, we review some concepts of dependence that are used in this paper. Readers who are not familiar with this area are advised to read Appendix A before reading the next two sections.

2. Bivariate IFR property

Recalling that a random variable X has IFR distribution if and only if $\mathcal{L}(X - t_1 | X > t_1) \geq_{\text{st}} \mathcal{L}(X - t_2 | X > t_2)$ for $t_1 < t_2$ and also if and only if $\mathcal{L}(X - t_1 | X > t_1) \geq_{\text{hr}} \mathcal{L}(X - t_2 | X > t_2)$ for $t_1 < t_2$, Bassan and Spizzichino (1999) considered the following two extensions of the concept of univariate IFR ageing to the bivariate case.

Definition 2.1. An exchangeable random vector $T = (T_1, T_2)$ is bivariate IFR (BIFR) if

$$\mathcal{L}(T_1 - t_1 \mid T_1 > t_1, T_2 > t_2) \geq_{st} \mathcal{L}(T_2 - t_2 \mid T_1 > t_1, T_2 > t_2) \quad \text{for } t_1 \leq t_2. \quad (2.1)$$

Definition 2.2. An exchangeable random vector (T_1, T_2) is said to be bivariate IFR in the strong sense (s-BIFR) if

$$\mathcal{L}(T_1 - t_1 \mid T_1 > t_1, T_2 > t_2) \geq_{hr} \mathcal{L}(T_2 - t_2 \mid T_1 > t_1, T_2 > t_2) \quad \text{for } t_1 \leq t_2. \quad (2.2)$$

Notice that (2.1) holds if and only if the joint survival function $\bar{F}(t_1, t_2) = P[T_1 > t_1, T_2 > t_2]$ of (T_1, T_2) is Schur-concave in (t_1, t_2) . For details on Schur-concavity, see Marshall and Olkin (1979); see also the discussion in Spizzichino (2001) and references therein for the use of Schur-concavity in multivariate ageing.

On the other hand, (T_1, T_2) is s-BIFR if and only if

$$R(t) = \frac{\bar{F}(x + t, y)}{\bar{F}(y + t, x)}$$

is increasing in t for $x < y$. Taking logs and then differentiating both sides, we see that this is equivalent to

$$r_{T_1 \mid T_2}(y + t \mid T_2 > x) \geq r_{T_1 \mid T_2}(x + t \mid T_2 > y) \quad \text{for } t > 0 \text{ and } 0 \leq x < y, \quad (2.3)$$

where $r_{T_1 \mid T_2}(\cdot \mid T_2 > t_2)$ denotes the hazard rate of the conditional distribution of T_1 given $\{T_2 > t_2\}$.

Unless the random variables T_1 and T_2 are independent, the two concepts of multivariate ageing as given by Definitions 2.1 and 2.2 are not equivalent. The latter implies the former, but the converse is not true; see the example presented in Remark 3.6 of Bassan and Spizzichino (1999), where some properties of these bivariate notions of ageing are studied in detail.

The first aspect that we focus on concerns the (unidimensional) ageing properties of the marginal distributions of the bivariate distributions under consideration. As was observed by Bassan and Spizzichino (1999), the marginals of a BIFR random vector need not be IFR unless the component lifetimes are independent. They show that, however, under some negative dependence between the components, the marginal distributions are NBU. In the next theorem we prove that under a suitable negative dependence condition between T_1 and T_2 , the marginals of s-BIFR random vectors are IFR.

Theorem 2.1. *Let (T_1, T_2) be s-BIFR and let T_1 be right tail decreasing in T_2 . Then the marginal distribution of T_1 is IFR.*

Proof. Since (T_1, T_2) being s-BIFR is equivalent to (2.3), taking $x = 0$ there, we get

$$\begin{aligned} r_{T_1}(y + t) &\geq r_{T_1 \mid T_2}(t \mid T_2 > y) \\ &\geq r_{T_1}(t) \end{aligned}$$

for $y > 0, t > 0$ by the RTD property. This proves the required result.

Now we give an example of an s-BIFR distribution which satisfies the conditions of the above theorem.

Example 2.1. Let the joint distribution of (T_1, T_2) be

$$\bar{F}(x, y) = \exp\{1 - \exp(x^2 + y^2)\}, \quad x \geq 0, y \geq 0.$$

It is easy to see that

$$R(t) = \frac{\bar{F}(x + t, y)}{\bar{F}(y + t, x)} = \exp\{e^{x^2+(y+t)^2} - e^{y^2+(x+t)^2}\}$$

is increasing in t for $x \leq y$. Hence (T_1, T_2) is s-BIFR. Also, T_1 is right tail decreasing in T_2 since

$$\frac{\bar{F}(t, y)}{\bar{F}(t, 0)} = \exp[e^{t^2} - e^{t^2+y^2}]$$

is clearly decreasing in t for $y > 0$. Hence, this bivariate distribution satisfies the conditions of the above theorem. It is s-BIFR and its marginals are also IFR.

An interesting fact in the literature and in applications of unidimensional ageing notions is that mixtures of distributions with some property of positive ageing do not necessarily maintain the same property (see e.g. the review paper of Shaked and Spizzichino (2001) and references therein; for a recent contribution, see Finkelstein and Esaulova (2001)). It is then of interest to find conditions on the mixtures under which one-dimensional or multivariate ageing properties are preserved.

We consider here conditionally i.i.d. pairs of lifetimes. In this case, both the joint law and the marginal law can be written as mixtures. It may be interesting to notice that bivariate ageing properties may be preserved under mixtures, whereas the opposite may hold for one-dimensional properties. For example, consider two i.i.d. IFR lifetimes. All the conditional joint laws are Schur-concave (BIFR), and hence so is the unconditional joint law, since Schur-concavity is preserved under mixtures (see Barlow and Mendel (1992) for a discussion on this issue). On the other hand, the marginal law is a mixture of IFR laws, and hence it need not be IFR.

In the next theorem, we give sufficient conditions under which the joint law of conditionally i.i.d. IFR random variables is s-BIFR. First, since we shall deal repeatedly with conditionally i.i.d. random variables, we single out the relevant assumptions and notations. Notice also that we will be repeatedly using the notion of TP_2 functions in this paper (see Karlin (1968)).

Hypotheses 2.1. Let Θ be a random variable taking values in a set L , Π its (prior) distribution and π its (prior) density (when it exists). Given that Θ, T_1 and T_2 are conditionally i.i.d. random variables with a common conditional survival function $\bar{G}(\cdot | \theta)$, the joint survival function is given by

$$\bar{F}(t_1, t_2) = \int_L \bar{G}(t_1 | \theta)\bar{G}(t_2 | \theta) d\Pi(\theta). \tag{2.4}$$

Theorem 2.2. Let Hypotheses 2.1 be satisfied, with $L \subset \mathbb{R}$. Assume also the following:

- (i) the mapping $(t, \theta) \mapsto \bar{G}(t | \theta)$ is log-concave (this implies, in particular, that all the conditional laws are IFR) and TP_2 ,
- (ii) the mixing distribution Π has a log-concave density π .

Then, for $t_1 < t_2$ and letting $D = \{T_1 > t_1, T_2 > t_2\}$, we have

$$\mathcal{L}(T_1 - t_1 | D) \geq_{hr} \mathcal{L}(T_2 - t_2 | D).$$

For the proof, we prefer to single out two simple lemmas which will be helpful in the next section as well.

Lemma 2.1. *Let $(x, z) \mapsto \phi(x, z)$ be (jointly) log-concave and TP_2 , $(x, y) \mapsto \psi(x, y)$ be TP_2 , $z \mapsto \psi(y, z)$ be log-concave for all y and $z \mapsto h(z)$ be log-concave. Then*

- (i) $(x, y) \mapsto \psi(x, y)h(y)$ is TP_2 ,
- (ii) $(x, z) \mapsto \phi(x, z)\psi(y, z)h(z)$ is log-concave for all y ,
- (iii) $(x, y) \mapsto \int \phi(x, z)\psi(y, z)h(z) dz$ is TP_2 ,
- (iv) $x \mapsto \int \phi(x, z)\psi(y, z)h(z) dz$ is log-concave for all y .

Proof. It is a simple matter to check that (i) and (ii) hold. Part (iii) follows from the basic composition formula (Karlin (1968)), and (iv) from Prékopa’s theorem (Prékopa (1973)).

Lemma 2.2. *Let $(x, y) \mapsto H(x, y)$ be TP_2 , and let $x \mapsto H(x, y)$ be log-concave for all y . Then, for $x < y$, the mapping*

$$t \mapsto R(t; x, y) := \frac{H(x + t, y)}{H(y + t, x)}$$

is increasing.

Proof. Write $R(t, x, y) = R_1(t, x, y)R_2(t, x, y)$, with

$$R_1(t, x, y) = \frac{H(x + t, y)}{H(y + t, y)}, \quad R_2(t, x, y) = \frac{H(y + t, y)}{H(y + t, x)}.$$

We will show that R_1 and R_2 are increasing in t . In fact, R_1 is increasing because $z \mapsto H(z, y)$ is log-concave for every y . Furthermore, from the TP_2 property of H we get

$$H(y + t + a, x)H(y + t, y) \leq H(y + t, x)H(y + t + a, y),$$

and it follows immediately that R_2 is increasing.

Proof of Theorem 2.2. Let $\phi(x, \theta) = \psi(x, \theta) = \bar{G}(x | \theta)$, let $h = \pi$ and let H be the joint survival function:

$$H(t_1, t_2) := \bar{F}(t_1, t_2) = \int \bar{G}(t_1 | \theta)\bar{G}(t_2 | \theta)\pi(\theta) d\theta.$$

We see that ϕ , ψ and h satisfy the conditions of Lemma 2.1, and hence H satisfies the conditions of Lemma 2.2. It follows that

$$t \mapsto \frac{\bar{F}(t_1 + t, t_2)}{\bar{F}(t_2 + t, t_1)}$$

is increasing, and the claim follows.

Remark 2.1. The conditions of Theorem 2.2 guarantee that the marginals are IFR, as can be easily checked using Prékopa’s theorem. These conditions are similar to those of Lynch (1999) for a mixture of IFR distributions to be IFR. In this respect we also notice the following: in

the case when T_1, T_2 are conditionally i.i.d. given Θ , the condition (2.2) can be rewritten in the form

$$\int_L \frac{\bar{G}(t_1 + r + \tau | \theta)}{\bar{G}(t_1 + r | \theta)} \pi(\theta | T_1 > t_1 + r, T_2 > t_2) d\theta$$

$$\geq \int_L \frac{\bar{G}(t_2 + r + \tau | \theta)}{\bar{G}(t_2 + r | \theta)} \pi(\theta | T_1 > t_1, T_2 > t_2 + r) d\theta \quad \text{for } t_2 > t_1 > 0 \text{ and } r, \tau > 0,$$

where we let

$$\pi(\theta | T_1 > t_1 + r, T_2 > t_2) \propto \pi(\theta) \bar{G}(t_1 + r | \theta) \bar{G}(t_2 | \theta),$$

which, by Bayes' formula, can be interpreted as the conditional distribution of Θ , given the observation of the survival data

$$\{T_1 > t_1 + r, T_2 > t_2\}.$$

An analogous result holds for $\pi(\theta | T_1 > t_1, T_2 > t_2 + r)$.

Notice, on the other hand, that the condition that the marginal distribution of T_1, T_2 be IFR can be rewritten in the form

$$\int_L \frac{\bar{G}(t_1 + \tau | \theta)}{\bar{G}(t_1 | \theta)} \pi(\theta | T_1 > t_1) d\theta \geq \int_L \frac{\bar{G}(t_2 + \tau | \theta)}{\bar{G}(t_2 | \theta)} \pi(\theta | T_1 > t_2) d\theta.$$

3. Bivariate DMRL property

In this section, we introduce a weak and a strong notion of bivariate DMRL ageing, in analogy with Definitions 2.1 and 2.2. A natural way to do so is to take into account the conditional mean residual life function given an observed bivariate survival data. A first, natural, definition in this vein is as follows.

Definition 3.1. An exchangeable random vector (T_1, T_2) is said to have bivariate DMRL (BDMRL) distribution if, for $t_1 < t_2$,

$$E[T_1 - t_1 | T_1 > t_1, T_2 > t_2] \geq E[T_2 - t_2 | T_1 > t_1, T_2 > t_2]. \tag{3.1}$$

Note that (T_1, T_2) is BDMRL if and only if one (and hence all) of the following equivalent conditions holds:

$$\int_{t_1}^{\infty} \bar{F}(x, t_2) dx \geq \int_{t_2}^{\infty} \bar{F}(x, t_1) dx \quad \text{for } t_1 < t_2,$$

$$\int_{t_1}^{\infty} \int_{t_2}^{\infty} \bar{F}(x, y) dx dy \text{ is Schur-concave in } (t_1, t_2),$$

$$\mu_{T_1 | T_2}(t_1 | T_2 > t_2) \geq \mu_{T_1 | T_2}(t_2 | T_2 > t_1) \quad \text{for } t_1 < t_2,$$

where $\mu_{T_1 | T_2}(\cdot | T_2 > t_2)$ denotes the mean residual life function of the conditional distribution of T_1 given $\{T_2 > t_2\}$.

Since stochastic ordering implies the ordering of the means, it follows immediately that the bivariate IFR property implies the bivariate DMRL property.

Note that for a univariate random variable X , the mean residual life $\mu_t(x)$ of X_t at time x satisfies the relation $\mu_t(x) = \mu_F(t + x)$.

Therefore,

$$\begin{aligned} X \text{ is DMRL} &\iff \mu_F(x) \text{ is decreasing in } x \\ &\iff \mu_t(x) \text{ is decreasing in } x \text{ for } t \geq 0, \\ &\iff \mathcal{L}(X - t_1 \mid X > t_1) \geq_{\text{mrl}} \mathcal{L}(X - t_2 \mid X > t_2) \text{ for } t_1 < t_2. \end{aligned}$$

Then, in the case of i.i.d. variables T_1, T_2 , the condition that they are DMRL is equivalent to

$$\mathcal{L}(T_1 - t_1 \mid T_1 > t_1, T_2 > t_2) \geq_{\text{mrl}} \mathcal{L}(T_2 - t_2 \mid T_1 > t_1, T_2 > t_2) \quad \text{for } t_1 < t_2. \quad (3.2)$$

Arguing as we did for the bivariate extension of the notion of IFR, notice now that the equivalence of (3.1) and (3.2) is no longer valid in the case when T_1, T_2 are exchangeable, but not independent, variables.

This encourages us to propose the following stronger notion of the bivariate DMRL property.

Definition 3.2. We say that an exchangeable random vector (T_1, T_2) is bivariate DMRL in the strong sense (s-BDMRL) if, for all $t_1 < t_2$, the inequality (3.2) holds.

Note that (T_1, T_2) is s-BDMRL if and only if one (and hence both) of the following equivalent conditions hold:

$$\begin{aligned} \mu_{T_1 \mid T_2}(x + t_1 \mid T_2 > t_2) \geq \mu_{T_1 \mid T_2}(x + t_2 \mid T_2 > t_1) \quad \text{for } t_1 < t_2 \text{ and for } x > 0, \\ \frac{\int_{x+t_1}^{\infty} \bar{F}(u, t_2) \, du}{\int_{x+t_2}^{\infty} \bar{F}(u, t_1) \, du} \text{ is increasing in } x > 0. \end{aligned}$$

In the following example, we give a bivariate law which is BDMRL but not s-BDMRL.

Example 3.1. Let

$$\bar{F}_{T_1, T_2}(u, v) \propto [1 + u^3 + v^3]^{-2}.$$

It is easy to check that this survival function is Schur-concave. Thus, the joint law is BIFR, and hence it is BDMRL. In order to see whether it is s-BDMRL, we have to determine whether

$$\frac{\int_{x+t_1}^{\infty} \bar{F}(u, t_2) \, du}{\int_{x+t_2}^y \bar{F}(u, t_1) \, du} = \frac{\int_{x+1}^{\infty} [65 + u^3]^{-2} \, du}{\int_{x+4}^{\infty} [2 + u^3]^{-2} \, du}$$

is increasing in x . On examining the plot of this function, we see that it is not monotone. Hence this distribution is not s-BDMRL. Since \geq_{hr} implies \geq_{mrl} , this distribution is not s-BIFR either.

In the next theorem, we give sufficient conditions for the bivariate s-DMRL property.

Theorem 3.1. Let

- (i) $\mu_{T_1 \mid T_2}(x \mid T_2 > y)$ be increasing in y for every $x > 0$, and
- (ii) $\mu_{T_1 \mid T_2}(x \mid T_2 > y)$ be decreasing in x for every $y > 0$ (that is, T_1 is conditionally DMRL for every given $T_2 > y$).

Then (T_1, T_2) is s-DMRL.

Proof. For $t_1 < t_2$ and $x > 0$,

$$\begin{aligned} \mu_{T_1 \mid T_2}(x + t_1 \mid T_2 > t_2) &\geq \mu_{T_1 \mid T_2}(x + t_2 \mid T_2 > t_2) \quad \text{(by (ii))} \\ &\geq \mu_{T_1 \mid T_2}(x + t_2 \mid T_2 > t_1) \quad \text{(by (i)).} \end{aligned}$$

This proves the required result.

In the next theorem, we show that under a certain type of negative dependence between the variables, the marginal distributions of a bivariate DMRL vector are NBUE.

Theorem 3.2. *Let (T_1, T_2) be bivariate DMRL and be negatively dependent in the sense that*

$$E(T_1 | T_2 > t) \leq E(T_1) \quad \text{for } t > 0. \tag{3.3}$$

Then the marginal distribution of T_1 is NBUE.

Proof. The vector (T_1, T_2) being DMRL is equivalent to

$$\mu_{T_1 | T_2}(t_2 | T_2 > t_1) \leq \mu_{T_1 | T_2}(t_1 | T_2 > t_2) \quad \text{for } t_1 < t_2.$$

Taking $t_1 = 0$, we get, for $t_2 > 0$,

$$\begin{aligned} \mu_{T_1}(t_2) &\leq \mu_{T_1 | T_2}(0 | T_2 > t_2) \\ &= E(T_1 | T_2 > t_2) \\ &\leq E(T_1). \end{aligned}$$

by (3.3). Hence T_1 is NBUE.

Remark 3.1. The condition (3.3) is implied by negative quadrant dependence between T_1 and T_2 .

In the next theorem, we give sufficient conditions under which the marginals of a multivariate s-DMRL distribution are DMRL.

Theorem 3.3. *Let (T_1, T_2) be s-DMRL and let*

$$\mu_{T_1 | T_2}(x | T_2 > y) \leq \mu_{T_1}(x) \quad \text{for } x, y > 0. \tag{3.4}$$

Then the marginal distribution of T_1 is DMRL.

Proof. The vector (T_1, T_2) being s-DMRL is equivalent to

$$\mu_{T_1 | T_2}(x + t_2 | T_2 > t_1) \leq \mu_{T_1 | T_2}(x + t_1 | T_2 > t_2) \quad \text{for } t_1 < t_2 \text{ and for } x > 0.$$

Taking $t_1 = 0$, we get

$$\mu_{T_1 | T_2}(x + t_2 | T_2 > 0) \leq \mu_{T_1 | T_2}(x | T_2 > t_2) \quad \text{for } t_1 < t_2 \text{ and for } x > 0.$$

That is, for $x \geq 0, t_2 \geq 0$,

$$\begin{aligned} \mu_{T_1}(x + t_2) &\leq \mu_{T_1 | T_2}(x | T_2 > t_2) \\ &\leq \mu_{T_1}(x) \end{aligned}$$

by (3.4), proving thereby that the distribution of T_1 is DMRL.

We now turn to consider the BDMRL and s-BDMRL properties for the case of conditionally i.i.d. variables. First, we show that conditionally i.i.d. DMRL random variables are bivariate DMRL.

Theorem 3.4. *Let T_1 and T_2 be conditionally i.i.d., as specified in Hypotheses 2.1. Let the conditional law of T_1 (and T_2) given $\Theta = \theta$ be DMRL for all θ . Then the joint distribution of (T_1, T_2) is bivariate DMRL.*

Proof. As seen earlier, conditionally on $\Theta = \theta$, T_1 is DMRL if and only if

$$\int_t^\infty \bar{G}(x | \theta) dx$$

is log-concave in t . Recalling (2.4), we have

$$\begin{aligned} \int_{t_1}^\infty \int_{t_2}^\infty \bar{F}(x, y) dx dy &= \int_{t_1}^\infty \int_{t_2}^\infty \left[\int_{\theta \in L} \bar{G}(x | \theta) \bar{G}(y | \theta) d\Pi(\theta) \right] dx dy \\ &= \int_{\theta \in L} \left[\int_{t_1}^\infty \int_{t_2}^\infty \bar{G}(x | \theta) \bar{G}(y | \theta) dx dy \right] d\Pi(\theta). \end{aligned}$$

Since this expression is a mixture of log-concave functions, it is Schur-concave. Hence the result.

In the next theorem we establish conditions under which conditionally i.i.d. DMRL random variables are bivariate DMRL in the strong sense.

Theorem 3.5. *Let T_1 and T_2 be conditionally i.i.d., as specified in Hypotheses 2.1, with $L \subset \mathbb{R}$ and Π admitting a density π . We assume the following:*

- (i) $\int_x^\infty \bar{G}(\xi | \theta) d\xi$ is jointly log-concave in θ and x (and hence, in particular, $\bar{G}(\cdot | \theta)$ is DMRL for all θ),
- (ii) $\bar{G}(v | \theta)$ is TP_2 and log-concave as a function of θ ,
- (iii) $\pi(\theta)$ is log-concave.

Then the joint distribution of (T_1, T_2) is bivariate s -DMRL.

Proof. Let \bar{F} be the joint survival function and let

$$H(t_1, t_2) := \int_{t_1}^\infty \bar{F}(u, t_2) du.$$

Clearly,

$$\begin{aligned} H(t_1, t_2) &= \int_{t_1}^\infty \left[\int_L \bar{G}(u | \theta) \bar{G}(t_2 | \theta) \pi(\theta) d\theta \right] du \\ &= \int_L \left[\int_{t_1}^\infty \bar{G}(u | \theta) du \right] \bar{G}(t_2 | \theta) \pi(\theta) d\theta. \end{aligned}$$

We need to show that, for $t_1 < t_2$,

$$t \mapsto \frac{H(t_1 + t, t_2)}{H(t_2 + t, t_1)}$$

is increasing.

Let

$$\phi(x, \theta) := \int_x^\infty \bar{G}(u | \theta) du, \quad \psi(x, \theta) = \bar{G}(x | \theta), \quad h = \pi.$$

Observe that $(t_1, \xi) \mapsto \mathbf{1}_{(t_1, \infty)}(\xi)$ is TP_2 , and hence by the basic composition formula ϕ is also TP_2 . Thus ϕ, ψ and h satisfy the conditions of Lemma 2.1, and hence H satisfies the conditions of Lemma 2.2 and the conclusion follows.

Appendix A. Some notions of dependence

There are several notions of positive and negative dependence between random variables and these have been discussed in detail in Barlow and Proschan (1981) and Shaked (1977). For a brief introduction, see Khaledi and Kochar (2000). The following concepts are used in this paper.

Let (T_1, T_2) be a bivariate random variable with joint density function $f(t_1, t_2)$ and joint survival function $\bar{F}(t_1, t_2) = P[T_1 > t_1, T_2 > t_2]$.

Definition A.1. The random variable (T_1, T_2) is said to be positively quadrant dependent (PQD) if

$$\{T_1 \mid T_2 > t_2\} \geq_{st} T_1 \quad \text{for all } t_2 > 0, \tag{A.1}$$

and negatively quadrant dependent (NQD) if the inequality in (A.1) is reversed.

Definition A.2. The random variable (T_1, T_2) is said to be right corner set increasing (RCSI) if

$$\begin{aligned} \{T_1 \mid T_2 > t_2\} &\geq_{hr} \{T_1 \mid T_2 > t_1\} \quad \text{for all } t_1 < t_2 & \tag{A.2} \\ \iff \bar{F}(t_1, t_2) &\text{ is TP}_2 \text{ in } (t_1, t_2) \\ \iff r_{T_1 \mid T_2}(t_1 \mid T_2 > t_2) &\text{ is decreasing in } t_2 \text{ for all } t_1 > 0. \end{aligned}$$

(Shaked (1977) calls this dependence DTP(1, 1).) If the inequality in (A.2) is reversed, we say that (T_1, T_2) is right corner set decreasing (RCSD).

Definition A.3. The random variable T_2 is said to be right tail increasing (RTI) in T_1 if

$$\begin{aligned} \{T_1 \mid T_2 > t_2\} &\geq_{hr} T_1 \quad \text{for all } t_2 > 0 & \tag{A.3} \\ \iff \frac{\bar{F}(t_1, t_2)}{\bar{F}(t_1)} &\text{ is increasing in } t_1 \text{ for all } t_2 > 0 \\ \iff P[T_2 > t_2 \mid T_1 > t_1] &\text{ is increasing in } t_1 \text{ for all } t_2 > 0. \end{aligned}$$

If the inequality in (A.3) is reversed, we say that T_2 is right tail decreasing (RTD) in T_1 .

Definition A.4. The random variables T_1, T_2 are dependent by total positivity of order (2, 1) (DTP(2, 1)) if

$$\begin{aligned} \int_t^\infty \bar{F}(x, y) dx &\text{ is TP}_2 \text{ in } (t, y) \\ \iff \mu_{T_1 \mid T_2}(t_1 \mid T_2 > t_2) &\text{ is increasing in } t_2 \text{ for all } t_1 > 0. \end{aligned}$$

If $\mu_{T_1 \mid T_2}(t_1 \mid T_2 > t_2)$ is decreasing in t_2 for all $t_1 > 0$, the variables are negatively dependent and T_1, T_2 are said to be DRR(2, 1).

We have the following implications among the above concepts of positive dependence:

$$DTP(2, 1) \iff RCSI \implies RTI \implies PQD.$$

Similar implications hold among the above concepts of negative dependence.

Acknowledgement

We thank an anonymous referee for useful suggestions.

References

- ARJAS, E. (1981). A stochastic process approach to multivariate reliability systems: notions based on conditional stochastic order. *Math. Operat. Res.* **6**, 263–276.
- BARLOW, R. E. AND MENDEL, M. B. (1992). De Finetti-type representations for life distributions. *J. Amer. Statist. Assoc.* **87**, 1116–1122.
- BARLOW, R. E. AND MENDEL, M. B. (1993). Similarity as a characteristic of ageing. In *Reliability and Decision Making*, eds R. E. Barlow, C. A. Clarotti and F. Spizzichino, Chapman and Hall, London, pp. 233–245.
- BARLOW, R. AND PROSCHAN, F. (1981). *Statistical Theory of Reliability and Life Testing*. To Begin With, Silver Spring, MD.
- BARLOW, R. E. AND SPIZZICHINO, F. (1993). Schur-concave survival functions and survival analysis. *J. Comput. Appl. Math.* **46**, 437–447.
- BASSAN, B. AND SPIZZICHINO, F. (1999). Stochastic comparison for residual lifetimes and Bayesian notions of multivariate ageing. *Adv. Appl. Prob.* **31**, 1078–1094.
- BRINDLEY, E. C. AND THOMPSON, W. A. (1972). Dependence and aging aspects of multivariate survival. *J. Amer. Statist. Assoc.* **67**, 822–830.
- DESHPANDE, J. V., KOCHAR, S. C. AND SINGH, H. (1986). Aspects of positive ageing. *J. Appl. Prob.* **23**, 748–58.
- FINKELSTEIN, M. S. AND ESAULOVA, V. (2001). Modeling a failure rate for a mixture of distribution functions. *Prob. Eng. Inf. Sci.* **15**, 383–400.
- GUESS, F. AND PROSCHAN, F. (1988). Mean residual life: theory and applications. In *Handbook of Statistics*, Vol. 7, *Quality Control and Reliability*, eds P. R. Krishnaiah and C. R. Rao, North-Holland, Amsterdam, pp. 215–224.
- KARLIN, S. (1968). *Total Positivity*. Stanford University Press.
- KHALEDI, B. AND KOCHAR, S. (2000). Stochastic comparisons and dependence among concomitants of order statistics. *J. Multivariate Anal.* **73**, 262–281.
- KOCHAR, S., MUKERJEE, H. AND SAMANIEGO, F. (2000). Estimation of a monotone mean residual life. *Ann. Statist.* **28**, 905–921.
- LYNCH, J. D. (1999). On conditions for mixtures of increasing failure rate distributions to have an increasing failure rate. *Prob. Eng. Inf. Sci.* **13**, 33–36.
- MARSHALL, A. W. AND OLKIN, I. (1979). *Inequalities: Theory of Majorization and Its Applications*. Academic Press, New York.
- PRÉKOPA, A. (1973). On logarithmic concave measures and functions. *Acta Sci. Math. (Szeged)* **34**, 335–343.
- SAVITS, T. H. (1985). A multivariate IFR distribution. *J. Appl. Prob.* **22**, 197–204.
- SHAKED, M. (1977). A family of concepts of dependence for bivariate distributions. *J. Amer. Statist. Assoc.* **72**, 642–650.
- SHAKED, M. AND SHANTHIKUMAR, J. G. (1988). Multivariate conditional hazard rates and the MIFRA and MIFR properties. *J. Appl. Prob.* **25**, 150–168.
- SHAKED, M. AND SHANTHIKUMAR, J. G. (1991). Dynamic multivariate aging notions in reliability theory. *Stoch. Process. Appl.* **38**, 85–97.
- SHAKED, M. AND SHANTHIKUMAR, J. G. (1994). *Stochastic Orders and Their Applications*. Academic Press, San Diego, CA.
- SHAKED, M. AND SPIZZICHINO, F. (2001). Mixtures and monotonicity of failure rate functions. In *Handbook of Statistics*, Vol. 20, *Advances in Reliability*, eds N. Balakrishnan and C. R. Rao, North-Holland, Amsterdam, pp. 185–198.
- SPIZZICHINO, F. (2001). *Subjective Probability Models for Lifetimes*. Chapman and Hall/CRC, Boca Raton, FL.