

A Primer on Laplacian Dynamics in Directed Graphs

J. J. P. Veerman* and R. Lyons†

(Dated: July 13, 2020)

We analyze the asymptotic behavior of general first order Laplacian processes on digraphs. The most important ones of these are diffusion and consensus with both continuous and discrete time. We treat diffusion and consensus as dual processes. This is the first complete exposition of this material in a single work.

I. INTRODUCTION

Directed graphs (or digraphs) are an important generalization of undirected graphs and they have wide-ranging applications. Examples include models of the internet [1] and social networks [2], food webs [3], epidemics [4], chemical reaction networks [5], databases [6], communication networks [7], the Pagerank algorithm [8], and networks of autonomous agents in control theory [9] to name but a few. In many of these applications, it is of crucial importance to understand the asymptotics (as $t \rightarrow \infty$) of solutions of the first order Laplacian differential equation

$$\dot{x} = -Lx \quad \text{and} \quad \dot{p} = -pL. \quad (\text{I.1})$$

We will show how the first of these equations is associated with the physical process of consensus and the second with diffusion. We will give a unified treatment of both of these in which one is treated as the dual to the other. We also show how this extends to the discrete versions of these processes.

We describe the basic theory of Laplacian dynamics on directed graphs that are weakly connected. The restriction of this theory to *undirected* graphs is well documented in textbooks (see [10], [11]), but as far as we know, this is the first complete exposition of the general theory (for *directed* graphs) in a single work.

Many of the results we will discuss had earlier been “folklore” results living largely outside the mathematics community and not always with complete proofs (see [12, 13] for some references). In the mathematics community, directed graphs are still much less studied than undirected graphs (especially true for the algebraic aspects). As a consequence, there are not many good mathematics books on the subject.

Part of the reason for that is probably that directed graphs are a lot messier than undirected graphs. For example, we will see that while *undirected* graphs are either connected or not, for *directed* graphs there are various gradations of connectedness. Another complication is that while Laplacians of *undirected* graphs are

diagonalizable and have real eigenvalues, neither statement is necessarily true for Laplacians of *digraphs*. Thus, many statements for undirected graphs take more work to prove, or are wrong.

Another reason for confusion is that there is no standard way to orient a graph. The in-degree Laplacian of G is the same as the out-degree Laplacian for G' , the graph with all orientations reversed. In [13], the convention was proposed where the direction of edges corresponds to the flow of information in the underlying problem. While here we are not discussing any particular applications, we can still make use of that convention. So in the case where there is a directed path from vertex i to vertex j , where will write that information goes from i to j , or, more succinctly, j “sees” i .

The set-up of this paper is as follows. In Section II we give the necessary definitions concerning directed graphs, and in Section III those concerning Laplacians. In Section IV, we discuss the spectrum of graph Laplacians, in particular the fact that all non-zero eigenvalues have positive real part. That means that the asymptotic behavior of the solutions of equations (I.1) is determined by the kernel of the Laplacians. Thus, in Section V we give a convenient basis for those eigenspaces. This allows us in Section VI to write the asymptotics in terms of that basis. This results in Theorem VI.1, which is perhaps the main result in this paper. In Section VII, we apply this to the most important of the Laplacians, namely the “random walk” Laplacian. This Laplacian is particularly suited to discretization of time, and we show that the asymptotics of the solution of the discretized equations is again essentially the same of that of the continuous time equations. We provide examples for all of our main statements.

Finally, a few notational issues. By “Laplacian” we mean a matrix of the form $E - ES$, where E is diagonal with positive entries on the diagonal and S is row stochastic (details are in Section III). Everything in this article goes through for matrices of the form $E - SE$. One only needs to exchange left and right eigenvectors. In the interest of brevity, we have not pursued this.

We use the notation $\mathbf{1}_S$ for the vector whose i th component is 1 if $i \in S$ and 0 elsewhere. That also means that $\mathbf{1}_{\{i\}}$ means the unit vector whose i th component equals 1 while being 0 everywhere else. The symbol e_i is used for the i th diagonal element of the matrix E .

These notes outline part of a series of 4 lectures given in summer-school/conference on mathematical modeling of complex systems in Pescara, 2019 [14]. Most of this

* Fariborz Maseeh Dept. of Math. and Stat., Portland State Univ.; veerman@pdx.edu

† Fariborz Maseeh Dept. of Math. and Stat., Portland State Univ.; rlyons@pdx.edu




FIG. II.1. This graph will serve as an example throughout the paper.

theory was described in [12, 13] and those are the two references that we rely most heavily on. However, various of those proofs have been substantially simplified, and other statements have been generalized (most notably Theorem V.2).

Acknowledgements: JJPV is grateful to the University of Chieti-Pescara for the generous hospitality offered.

II. GRAPH THEORETIC DEFINITIONS

Definition II.1 A directed graph (or digraph) is a set $V = \{1, \dots, n\}$ of vertices together with a set $E \subseteq V \times V$ or ordered pairs (the edges).

The graph in Figure II.1 will serve as our example of a digraph. Edges will be indicated by $i \rightarrow j$ or (i, j) (or ij for short). So the graph in the figure has edges $(1, 2)$, $(1, 6)$, $(6, 7)$, et cetera, but it does *not* have the edges $(2, 1)$ and $(6, 1)$. Directed paths from i to j are denoted by $i \rightsquigarrow j$. We will express this informally as: information goes from i to j , or: j “sees” i . For example, the graph in Figure II.1 has a path $4 \rightsquigarrow 6$, but there is no path $6 \rightsquigarrow 4$.

Connectedness for *undirected* graph is straightforward: an undirected graph is either connected or it is not. However, for a digraph, the situation is slightly more complicated. We need the notion of *underlying graph*. This is the undirected graph one obtains by erasing the direction of the edge. Equivalently, it is the graph obtained by adding to each directed edge an edge in the opposite direction.

Definition II.2 i) A digraph G is strongly connected if for every ordered pair of vertices (i, j) , there is a path $i \rightsquigarrow j$. Equivalently, if for every pair i and j : $i \rightsquigarrow j$.
 ii) A digraph G is unilaterally connected if for every ordered pair of vertices (i, j) , there is a path $i \rightsquigarrow j$ or a path $j \rightsquigarrow i$.
 iii) A digraph G is weakly connected if the underlying undirected graph is connected.
 iv) A digraph G is not connected if it is not weakly connected.

A subgraph which is strongly connected is called a

strongly connected component. We will frequently abbreviate this to SCC.

The study of a graph that is not connected is of course equivalent to the study of each its components. So the most general graph we want to study is weakly connected. The graph of Figure II.1 is an example of such a graph. We will need some terminology to indicate certain subgraphs. We borrow our terminology from [12] and [13].

Definition II.3 i) Let $i \in V$. The reachable set $R(i)$ consists of all $j \in V$ with $i \rightsquigarrow j$.
 ii) A reach R is a maximal reachable set, or a maximal unilaterally connected set.
 iii) A cabal $B \subseteq R$ is the set of vertices from which the entire reach R is reachable. If it is a single vertex, it is usually called a leader or a root.
 iv) The exclusive part $H \subseteq R$ are those vertices in R that do not “see” vertices from other reaches.
 v) The common part $C \subseteq R$ are those vertices in R that also “see” vertices from other reaches.

Note that every reach has a single non-empty cabal. We illustrate these ideas using the graph in Figure II.1. That graph has two reaches, $R_1 = \{1, 2, 6, 7\}$ and $R_2 = \{3, 4, 5, 6, 7\}$. Their exclusive parts are $H_1 = \{1, 2\}$ and $H_2 = \{3, 4, 5\}$. The common parts are $C_1 = C_2 = \{6, 7\}$. Finally, the cabals are $B_1 = \{1\}$ and $B_2 = \{3, 4, 5\}$. It is an interesting exercise to reverse the orientation of the edges and do the taxonomy again. It is easy to see that the graph has again two reaches. But in general the number of reaches need not be constant under orientation. Consider for example the graph $1 \rightarrow 2 \leftarrow 3$.

The relation $i \rightsquigarrow j$ between vertices of G that defines an SCC (see Definition II.2) an equivalence relation. Thus it gives a unique partition of the vertices of G .

Definition II.4 The condensation $SC[G]$ of G is the graph obtained by identifying vertices of the same SCC (or grouping them together). See Figure II.2.




FIG. II.2. The condensation $SC[G]$ of G in Figure II.1.

These equivalence classes respect the categories of Definition II.3. For example, given that i is in a cabal, then

$i \leftrightarrow j$ is equivalent to j is in the same cabal. We leave it to the reader to check the other categories. Notice that $SC[G]$ can have no cycles and therefore all the cabals are singletons.

Definition II.5 Given a digraph G with vertex i , then N_i stands for the set of vertices j such that there is an edge $j \rightarrow i$. This is also called the (in-degree) neighborhood of i .

III. GRAPH LAPLACIANS

Definition III.1 The combinatorial adjacency matrix Q of the graph G is defined as $Q_{ij} > 0$ if there is an edge ji (if “ i sees j ”) and 0 otherwise. If vertex i has no incoming edges, set $Q_{ii} = 1$ (create a loop).

The last convention, on loops, is only adopted to ensure that the degree matrix D , defined below, can be taken to be non-singular (and thus invertible). One can drop the convention, but then one has to define the so-called pseudo-inverse of D . This is the approach taken in [12]. The two approaches are equivalent.

The non-zero values of Q_{ij} are the *weights* of the edges (j, i) . In the interest of brevity, our main example in Figure II.1 has unit weights. However, everything goes through in the general case.

Definition III.2 The in-degree matrix D is a diagonal matrix whose diagonal entry corresponding to the vertex i equals the sum of the weights of the edges ji arriving at i : $d_i \equiv \sum_j Q_{ij}$.

The matrices D and Q are used to generate S , a row stochastic (non-negative, every row adds to 1) version of the adjacency matrix.

Definition III.3 The (row) stochastic matrix $S \equiv D^{-1}Q$ is called the normalized adjacency matrix.

Definition III.4 Let E be a non-negative diagonal matrix. A Laplacian is a matrix of the form $E - ES$. Common examples are: the combinatorial (comb) Laplacian, $L^c \equiv D - DS = D - Q$, and the random walk (rw) Laplacian, $\mathcal{L} \equiv I - S$.

Here Q , D , and S are as defined earlier.

If L is a Laplacian matrix, then $(Lx)_k$ is equal to $\sum_{i \neq k} \alpha_i (x_k - x_i)$ for some combination of α_i . Thus, Laplacians describe relative observations. This is usually called “decentralized”. Clearly, matrices with this property must have row sum zero. It shares this property — and hence the name — with the discretization of the second derivative (or “Laplacian”) of a function $f : \mathbb{R} \rightarrow \mathbb{R}$:

$$f''(j) \approx f(j-1) - 2f(j) + f(j+1)$$

Notice, however, that in Definition III.4, the above expression would be the negative of a Laplacian. This convention we use ensures that Laplacians have eigenvalues whose real part is non-negative.

As an example, we work out the matrices corresponding to the graph G of Figure II.1 assuming all weights are 1.

$$Q = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{pmatrix} \quad D = \text{diag} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 2 \\ 2 \end{pmatrix} \quad (\text{III.2})$$

$$L^c \equiv D - Q = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 \\ -1 & 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & -1 & 0 & 0 & -1 \end{pmatrix} \quad (\text{III.3})$$

The spectrum of L^c is:

$$\left\{ 0, 0, 1, 1, 3, \frac{3}{2} + i \frac{\sqrt{3}}{2}, \frac{3}{2} - i \frac{\sqrt{3}}{2} \right\}. \quad (\text{III.4})$$

The random walk Laplacian $\mathcal{L} \equiv I - D^{-1}Q$ is,

$$\mathcal{L} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 \\ -1/2 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & -1/2 & 0 & 0 & -1/2 \end{pmatrix} \quad (\text{III.5})$$

The spectrum of \mathcal{L} is given by:

$$\left\{ 0, 0, \frac{1}{2}, 1, \frac{3}{2}, \frac{3}{2} + i \frac{\sqrt{3}}{2}, \frac{3}{2} - i \frac{\sqrt{3}}{2} \right\}. \quad (\text{III.6})$$

As this example shows the Laplacians do not necessarily have a real spectrum. Nor, in fact, do they necessarily have a complete basis of eigenvectors. We leave it to the reader to verify that in Figure III.3, $L[G_1]$ and $\mathcal{L}[G_1]$ (with all weights equal to 1) have a non-real spectrum and that $\mathcal{L}[G_2]$ and $L[G_3]$ have a non-trivial Jordan block of dimension 2.

Definition III.5 Given a graph G , with $S = D^{-1}Q$. Let E^+ and E be non-negative diagonal matrices such





FIG. III.3. From left to right, top to bottom, three strongly connected graphs G_1 , G_2 , and G_3 .

that $E^+ \geq E$ (entry-wise). A generalized Laplacian is a matrix of the form $M = E^+ - ES$. The matrix is strict generalized if $E^+ \neq E$. Common examples are: $M^c = E^+ - DS$ with $E = D$ (comb), and $M = I - ES$ with $E^+ = I$ (rw).

For us, the importance of this definition lies in the fact that the characteristic polynomial of the Laplacian of a digraph is a product of characteristic polynomials of generalized Laplacians (see Proposition IV.3). For example, in equation (III.3), the diagonal blocks are generalized comb Laplacians, and in (III.5), they are generalized rw Laplacians.

IV. SPECTRA OF GRAPH LAPLACIANS

Lemma IV.1 *Let G be an undirected graph. The eigenvalues of a generalized Laplacian M are real and the eigenvectors form a complete basis. Neither holds necessarily even for Laplacians of a strongly connected digraph.*

Proof. A matrix that is conjugate to a real symmetric matrix has real eigenvalues and its eigenvectors form a complete basis. Now, recalling that $S = D^{-1}Q$ and that Q is symmetric because G is undirected, we set $H := ED^{-1}$, a diagonal matrix with positive diagonal,

and derive

$$\begin{aligned} M &= E^+ - ES = E^+ - HQ \\ &= E^+ - H^{1/2}H^{1/2}QH^{1/2}H^{-1/2} \\ &= H^{1/2} \left(E^+ - H^{1/2}QH^{1/2} \right) H^{-1/2}. \end{aligned}$$

The last equality holds, because diagonal matrices commute. Since $E^+ - H^{1/2}QH^{1/2}$ is symmetric, M is conjugate to a symmetric matrix.

The counter-examples of Section III, Figure III.3, establish the second part of this lemma. ■

Proposition IV.2 *Every non-zero eigenvalue of a generalized Laplacian has positive real part.*

Proof. Denote the diagonal elements of E^+ and E by e_i^+ and e_i , respectively. We have $e_i^+ \geq e_i > 0$. Apply Gersgorin's theorem [15] to

$$(E^+ - ES)v = \lambda v.$$

It follows that all eigenvalues are in the union of the closed balls

$$B_{e_i}(e_i^+) = \{x \in \mathbb{C} \mid |x - e_i^+| \leq e_i\}.$$

The statement follows. ■

Proposition IV.3 *The adjacency matrix of $SC[G]$ is lower block triangular after a reordering of the vertices of G .*

Proof. $SC[G]$ (see Definition II.4) cannot contain any cycles because the SCC's represented by vertices in a cycle of $SC[G]$ would in fact form a larger SCC. The graph associated with $SC[G]$ can be drawn with the arrows pointing upward (see Figure II.2). Then its vertices can be relabeled so that the vertex at the upper end (head) of an edge is greater than the vertex at its tail. This is equivalent to saying that $SC[G]$ is lower block triangular. ■

Corollary IV.4 *Any generalized Laplacian $M = E^+ - ES$ is lower block triangular after a reordering of the vertices of G . The characteristic polynomial of M is the product of the characteristic polynomials of the $M[K_i]$ where the K_i are the SCC's of G .*

Proposition IV.5 *Let G be an SCC. Any strict generalized Laplacian M is non-singular. Any Laplacian L has eigenvalue 0 with geometric and algebraic multiplicity 1.*

Proof. Let $M = E^+ - ES$ and suppose it has an eigenpair $\{0, v\}$. We can renormalize v so that the component with the largest modulus is $v_k = 1$. $Mv = 0$ gives:

$$\frac{e_k^+}{e_k} = \sum_{j \in \mathcal{N}_k} S_{kj} v_j,$$

where \mathcal{N}_k stands for the *neighborhood* of k (see Definition II.5). The left hand of this equality is greater than or equal to 1. The right hand is an *average* over entries with modulus less than or equal to 1. The only way the sum can equal 1 is if $v_j = 1$ for all $j \in \mathcal{N}_k$. Thus

$$e_k^+ = e_k \quad \text{and} \quad \forall j \in \mathcal{N}_k : v_j = 1. \quad (\text{IV.7})$$

Given any vertex i , there is a path $i \rightsquigarrow k$ (G is SCC), and thus (IV.7) holds for any vertex i .

The above reasoning proves that M has eigenvalue 0 if and only if it is an actual Laplacian (i.e. $E^+ = E$). Furthermore, it shows that all members of the kernel of an actual Laplacian L are multiples of $\mathbf{1}$ (the all ones vector). It remains to show that, in the case of a Laplacian $L = M$, the algebraic multiplicity of the eigenvalues 0 equals 1.

If 0 has algebraic multiplicity $m > 1$, there is a vector x such that

$$L^m x = 0 \quad \text{and} \quad L^{m-1} x \neq 0.$$

This means that $L^{m-1} x = L(L^{m-2} x) = \mathbf{1}$. So $v = L^{m-2} x$ has $Lv = \mathbf{1}$. Suppose that $\text{Re}(v_i)$ is minimized at $i = k$. Then $Lv = \mathbf{1}$ leads to

$$e_k \text{Re}(v_k) = 1 + e_k \sum_j S_{kj} \text{Re}(v_j) \geq 1 + e_k \text{Re}(v_k),$$

which is a contradiction. ■

Remark: An alternative proof is possible using the Perron Frobenius theorem [8, 16] to solve $Sv = v$. For this, one must first note that being an SCC means that the normalized adjacency matrix S is irreducible.

Theorem IV.6 *Given a digraph G . The algebraic and geometric multiplicity of the eigenvalue 0 of L equals the number of reaches. A strict generalized Laplacian is non-singular. All non-zero eigenvalues have positive real part.*

Proof. We can partition the vertices of G in SCC's. By Corollary IV.4, upon reshuffling the SCC's, the resulting Laplacian matrix is lower block triangular, and each diagonal block is a generalized Laplacian. By Proposition IV.5, the geometric and algebraic multiplicity of 0 equals the number of diagonal blocks (or SCC's) that are *actual* Laplacians. The generalized Laplacian of a diagonal is an actual Laplacian if and only if that SCC has no edges coming in from other SCC's. But that happens if and only if that SCC is a cabal. The number of cabals equals the number of reaches. ■

V. KERNELS RIGHT AND LEFT

Theorem V.1 *Let G be a digraph with $k \geq 1$ reaches. The right kernel of a Laplacian L consists of the column vectors $\{\gamma_1, \dots, \gamma_k\}$, where:*

$$\begin{cases} \gamma_{m,j} = 1 & \text{if } j \in H_m \quad (\text{exclusive}) \\ \gamma_{m,j} \in (0, 1) & \text{if } j \in C_m \quad (\text{common}) \\ \gamma_{m,j} = 0 & \text{if } j \notin R_m \quad (\text{not in reach}) \\ \sum_{m=1}^k \gamma_{m,j} = \mathbf{1} \end{cases}$$

Proof. Pick any of the k reaches and denote it by R . Denote its exclusive respectively, common parts by H and C . Recall (after Definition II.4) that the SCC's respect these categories. Thus, let X consist of the SCC's outside R that are "seen" by C , and Z the ones that are not "seen" by the cabals in C .

We can chop up the Laplacian by looking at the interactions between those four groupings of SCC's: H , X , C , and Z . For example, X does not "see" Z , because otherwise C would also "see" Z . Similarly, X does not "see", because otherwise X would "see" H , and therefore be part of the same reach. This way, we obtain the schematic Laplacian given in equation (V.8). This matrix is block triangular in agreement with Proposition IV.3.

We obtain a vector in the null space of L if we can solve the following equation.

$$\begin{pmatrix} L_{HH} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & L_{XX} & \mathbf{0} & \mathbf{0} \\ L_{CH} & L_{CX} & L_{CC} & \mathbf{0} \\ \mathbf{0} & L_{ZX} & \mathbf{0} & L_{ZZ} \end{pmatrix} \begin{pmatrix} \mathbf{1}_H \\ \mathbf{0}_X \\ \mathbf{x}_C \\ \mathbf{0}_Z \end{pmatrix} = \begin{pmatrix} \mathbf{0}_H \\ \mathbf{0}_X \\ \mathbf{0}_C \\ \mathbf{0}_Z \end{pmatrix}. \quad (\text{V.8})$$

But this equation boils down to

$$L_{HH} \mathbf{1}_H = \mathbf{0}_H \quad \text{and} \quad L_{CH} \mathbf{1}_H + L_{CC} \mathbf{x}_C = \mathbf{0}_C.$$

The first of these is satisfied since L is a Laplacian (row-sum zero) and therefore so is L_{HH} . The second of these has a unique, real solution if L_{CC} is invertible. The latter is true, because we can partition C into SCC's K_i in such a way that L_{CC} becomes lower block triangular (Proposition IV.3) and the restriction of L_{CC} to a block K_i is a strict generalized Laplacian and so has strictly positive eigenvalues (Proposition IV.5). Thus L_{CC} is non-singular (Corollary IV.4).

Denote this real eigenvector with eigenvalue 0 by v . Suppose that the maximum component is v_n . Then the same reasoning that leads to (IV.7) shows that if n "sees" a vertex j , then $v_j = v_n$. Thus, since n "sees" the cabal where $v_i = 1$, we must have $v_n = 1$. Similarly, by supposing that v_n is the minimum component of v_i , see that $v_n = 0$. Thus all values of v_i are in $[0, 1]$.

Every vertex i in C "sees" a vertex in H (with value 1) and a vertex in X (with value 0). Thus the value at i is ultimately an average collection of values that contain both 0 and 1. Thus all entries (in C) are in $(0, 1)$.

Finally, $\sum_{m=1}^k \gamma_{m,j} = \mathbf{1}$, because $L\mathbf{1}$ must be zero, and $\sum_{m=1}^k \gamma_{m,j}$ is the only combination of the γ 's that equals 1 on every vertex of the exclusive parts. ■

As an example, we compute the basis of the null space for the Laplacian given in (III.3) or (III.5) (they have the same null space).

$$\begin{aligned} \gamma_1 &= \left(1 \ 1 \ 0 \ 0 \ 0 \ \frac{2}{3} \ \frac{1}{3} \right)^T \quad \text{and} \\ \gamma_2 &= \left(0 \ 0 \ 1 \ 1 \ 1 \ \frac{1}{3} \ \frac{2}{3} \right)^T \end{aligned}$$

We now study the *left kernel* of L . As a mnemonic, we use the following device: the horizontal “overbar” on a vector $\bar{\gamma}$ indicates a (horizontal) row vector.

Theorem V.2 *Let G be a digraph with $k \geq 1$ reaches. The left kernel of Laplacian L consists of the row vectors $\{\bar{\gamma}_1, \dots, \bar{\gamma}_k\}$, where:*

$$\left\{ \begin{array}{ll} \bar{\gamma}_{m,j} > 0 & \text{if } j \in B_m \quad (\text{cabal}) \\ \bar{\gamma}_{m,j} = 0 & \text{if } j \notin B_m \quad (\text{not in cabal}) \\ \sum_{j=1}^k \bar{\gamma}_{m,j} = 1 \\ \{\bar{\gamma}_m\}_{m=1}^k \text{ are orthogonal} \end{array} \right.$$

Proof. The geometric and algebraic multiplicities of the eigenvalue 0 of L equal k (Theorem IV.6). All we have to do is: find k vectors $\bar{\gamma}_i$ in the left kernel of $L = E(I - S)$.

For each reach R , we split the vertices into the cabal B of R and the “rest”, X . We obtain an vector in the left null space of L if we can solve the following equation.

$$\left(\bar{\mathbf{x}}_B \ \bar{\mathbf{0}}_X \right) \begin{pmatrix} L_{BB} & \mathbf{0} \\ L_{XB} & L_{XX} \end{pmatrix} = \left(\bar{\mathbf{0}}_B \ \bar{\mathbf{0}}_X \right). \quad (\text{V.9})$$

B is an SCC and so by Proposition IV.5, this has a unique solution of the form $\bar{\gamma} = (\bar{\gamma}_B, \bar{\mathbf{0}}_X)$.

Set $\bar{v}_B = \bar{\gamma}_B E^{-1}$. Then \bar{v}_B satisfies

$$\bar{v}_B (I_{BB} - S_{BB}) = 0,$$

where S_{BB} is row stochastic and irreducible. The positivity of v_B follows from Perron Frobenius. (A direct proof would take a little longer.) Thus $\bar{\gamma}_B$ is strictly positive. Items ii and iv of the theorem follow after normalizing. ■

The left null spaces for the Laplacians in equations (III.3) and (III.5) are the same and are spanned by:

$$\begin{aligned} \bar{\gamma}_1 &= \left(1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \right) \quad \text{and} \\ \bar{\gamma}_2 &= \left(0 \ 0 \ \frac{1}{3} \ \frac{1}{3} \ \frac{1}{3} \ 0 \ 0 \right). \end{aligned}$$

For future reference, we include this definition.

Definition V.3 *For a digraph G with n vertices with k reaches, we define the $n \times n$ matrix Γ whose entries are given by:*

$$\Gamma_{ij} \equiv \sum_{m=1}^k \gamma_{m,i} \bar{\gamma}_{m,j} \quad \text{or} \quad \Gamma = \sum_{m=1}^k \gamma_m \otimes \bar{\gamma}_m$$

For the Laplacians given in (III.3) and (III.5), we obtain

$$\Gamma = \sum_{m=1}^k \gamma_m \otimes \bar{\gamma}_m = \frac{1}{9} \begin{pmatrix} 9 & 0 & 0 & 0 & 0 & 0 & 0 \\ 9 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 3 & 3 & 0 & 0 \\ 0 & 0 & 3 & 3 & 3 & 0 & 0 \\ 0 & 0 & 3 & 3 & 3 & 0 & 0 \\ 6 & 0 & 1 & 1 & 1 & 0 & 0 \\ 3 & 0 & 2 & 2 & 2 & 0 & 0 \end{pmatrix}. \quad (\text{V.10})$$

Lemma V.4 *Let L be an $n \times n$ Laplacian matrix. The kernels of Theorems V.1 and V.2 can be extended to bases of (generalized) right eigenvectors (columns), $\{\gamma_i\}_{i=1}^n$, and of (generalized) left eigenvectors (rows), $\{\bar{\gamma}_i\}_{i=1}^n$, such that the matrices:*

$$H = \left(\gamma_1 \ \gamma_2 \ \dots \ \gamma_n \right) \quad \text{and} \quad \bar{H} = \begin{pmatrix} \bar{\gamma}_1 \\ \bar{\gamma}_2 \\ \vdots \\ \bar{\gamma}_n \end{pmatrix}$$

are inverses of one another.

Proof: The right and left eigenvectors of L defined in Theorems V.1 and V.2 already satisfy $\bar{\gamma}_i \gamma_j = \delta_{ij}$ for $i, j \in \{1, \dots, k\}$.

The extension follows directly from the Jordan Decomposition Theorem [15]. Let J be the Jordan normal form of L . Then that theorem tells us that there is an invertible matrix H such that $LH = HJ$ or $H^{-1}L = JH^{-1}$. Right multiply the first equation by the standard column basis vector $\mathbf{1}_{\{i\}}$ to show that the i th column of H is a generalized right eigenvector. Left multiply by $\mathbf{1}_{\{i\}}^T$ to see that the i th row of H^{-1} is a generalized left eigenvector. ■

Definition V.5 *Let $\{\gamma_i\}_{i=1}^k$ be the (column) vectors of Theorem V.1 and $\{\bar{\gamma}_i\}_{i=1}^k$ the (row) vectors of Theorem V.2. Define*

$$H^0 = \left(\gamma_1 \ \gamma_2 \ \dots \ \gamma_k \right) \quad \text{and} \quad \bar{H}^0 = \begin{pmatrix} \bar{\gamma}_1 \\ \bar{\gamma}_2 \\ \vdots \\ \bar{\gamma}_k \end{pmatrix}.$$

From Definition V.3, we now easily compute the following.

Lemma V.6 $H^0 \bar{H}^0 = \Gamma$.

Notice the difference between Lemmas V.4 and V.6. The matrices H and \bar{H} are both $n \times n$ so that $\bar{H}H = I$ implies $H\bar{H} = I$. However, H^0 is a $n \times k$ matrix and \bar{H}^0 is $k \times n$ and there is no such simplification.

VI. LAPLACIAN DYNAMICS: CONSENSUS AND DIFFUSION

Throughout this section, we will assume that the digraph G has k reaches and that L is a Laplacian of G . In what follows x will always stand for a column vector and p for a row vector. In this section we are interested in solving the first order Laplacian equations:

$$\begin{aligned} \dot{x} &= -Lx \text{ with } x(0) = x_0, \\ \dot{p} &= -pL \text{ with } p(0) = p_0. \end{aligned} \quad (\text{VI.11})$$

The first of these equations is usually called *consensus* and the second is its dual problem of *diffusion*. We shall see below why these names are appropriate and how the solutions of these two problems are related. We start by discussing the solutions to the consensus problem.

Theorem VI.1 $\lim_{t \rightarrow \infty} e^{-Lt} = \Gamma$.

Proof: Let $\{\gamma_i\}_{i=1}^k$ and $\{\bar{\gamma}_i\}_{i=1}^k$ as in Theorems V.1 and V.2, and then extends these sets to complete basis of (generalized) eigenvectors $\{\gamma_i\}_{i=1}^n$ and $\{\bar{\gamma}_i\}_{i=1}^n$ as in Lemma V.4. Let λ_i be the eigenvalue associated with the i th (generalized) eigenvector γ_i (or $\bar{\gamma}_i$). So an initial condition $x_0 = y_0 + z_0$ can be decomposed as

$$\begin{aligned} x_0 &= y_0 + z_0 \\ \text{where } y_0 &= \sum_{i=1}^k \alpha_i \gamma_i \\ \text{and } z_0 &= \sum_{i=k+1}^n \alpha_i \gamma_i. \end{aligned} \quad (\text{VI.12})$$

From the standard theory of linear differential equations (see, for example, [17]), one easily derives that the general solution of the consensus problem $\dot{x} = -Lx$ is given by

$$x(t) = e^{-Lt} x_0 = \sum_{i=1}^n \gamma_i e^{-\lambda_i t} \xi_i(t), \quad (\text{VI.13})$$

where $\xi_i(t)$ are polynomials whose degrees are less than the size of the Jordan block corresponding to λ_i . Furthermore, if the dimension of that Jordan block equals 1, then $\xi_i(t) = \alpha_i$. By Theorem V.1, we have $\lambda_i = 0$ for $i \in \{1, \dots, k\}$. Also the zero eigenvalue has only trivial Jordan blocks and so for $i \in \{1, \dots, k\}$, $\xi_i = 1$ and $\beta_i = \alpha_i$. By Theorem IV.6, the λ_i , in terms with $i > k$, have positive real parts, and so these terms converge to

zero. Therefore, substitute equation (VI.12) into equation (VI.13) to get

$$\lim_{t \rightarrow \infty} x(t) = \sum_{m=1}^k \alpha_m \gamma_m = y_0.$$

Next, we determine the α_i . Definition V.5 and Lemma V.4 imply that

$$\begin{aligned} H^0 \bar{H}^0 x_0 &= H^0 \left(\bar{H}^0 \sum_{i=1}^n \alpha_i \gamma_i \right) \\ &= H^0 \left(\sum_{i=1}^k \alpha_i \mathbf{1}_{\{i\}} \right) = \sum_{i=1}^k \alpha_i \gamma_i = y_0. \end{aligned}$$

Note the change in the upper limit in the middle equality. Notice also that the vector $\mathbf{1}_{\{i\}}$ is a k -dimensional column vector, as opposed to γ_i which is n -dimensional. The result follows from Lemma V.6. ■

Since the the solutions of equation (VI.11) are given by $e^{-Lt}x(0)$ and $p(0)e^{-Lt}$, we have the following corollary.

Corollary VI.2 *The solutions of (VI.11) satisfy:*

$$\lim_{t \rightarrow \infty} x(t) = \Gamma x_0 \quad \text{and} \quad \lim_{t \rightarrow \infty} p(t) = p_0 \Gamma$$

As an example, let us consider the equations (VI.11) for the comb Laplacian L^c of Figure II.1 with initial conditions x_0 and p_0 concentrated on vertex 7 only. Then, from (V.10), we get

$$\begin{aligned} \lim_{t \rightarrow \infty} x(t) &= \Gamma x_0 = \mathbf{0} \quad \text{and} \\ \lim_{t \rightarrow \infty} p(t) &= p_0 \Gamma = \frac{1}{9}(3, 0, 2, 2, 2, 0, 0). \end{aligned}$$

We return to the first part of equation (VI.11) to explain why this called the consensus problem. The row sum of the Laplacian is zero, so we have $\mathcal{L}\mathbf{1} = 0$. Thus $\mathbf{1}$ is in the right kernel of L . If the eigenvalue 0 is non-degenerate, then from Corollary VI.2, we conclude that $\mathbf{1}$ is the final state and every component of the vector x has the same value. The system is, as it were, in complete agreement or consensus. Write out the differential equation in more detail and you get

$$\dot{x}_i = \sum_j e_i S_{ij} (x_j - x_i).$$

Thus \dot{x}_i is influenced by the *relative* (to x_i itself) positions of x_j where ji is a directed edge. In terms of Definition II.5, j is in the (in-degree) neighborhood of i . In other words, the consensus flows in the *same* direction as the information. In our formulation, consensus describes how the information spreads over the whole graph. Another way of saying this is that the influence of a vertex i exercises over the other vertices is described by $\Gamma \mathbf{1}_{\{i\}}$.

Next we explain how do these theorems apply to the diffusion problem in the second part of equation (VI.11). In this case, the vanishing of the row sum of the Laplacian implies that $\sum_i \dot{p}_i = 0$. Thus the sum of the components of p is preserved. Writing out the equation in full, we get

$$\dot{p}_i = \sum_j p_j e_j S_{ji} - p_i e_i. \quad (\text{VI.14})$$

Thus, if all p_j are non-negative and $p_i = 0$, then $\dot{p}_i > 0$. This means that the the positive p orthant is preserved. Since probability (or mass) is non-negative, these two observations together mean that the second process preserves total probability or mass. Hence the name diffusion. It is important to note that (VI.14) implies that p_i is influenced by the strengths of p_j where ij is a directed edge. In terms of Definition II.5, i is in the (in-degree) neighborhood of j . In other words, diffusion flows in the direction *contrary* to the direction of the information. Diffusion, in our formulation, tracks the source of the information. Another way of saying this is that the influencers of a vertex i are described by $\mathbf{1}_{\{i\}}^T \Gamma$.

VII. THE DISCRETIZATION OF THE RW LAPLACIAN

Perhaps the most important example of Laplacians is the rw Laplacian \mathcal{L} of Definition III.4 (with or without weighting the edges). It is most frequently used, among other things, to describe diffusion and consensus related problems, in discrete as well as continuous time. In contrast with the more general Laplacian, it is particularly well-behaved if it is discretized using time step 1. For the discrete consensus problem, we obtain

$$\begin{aligned} x(\ell + 1) - x(\ell) &= (S - I)x(\ell) \\ \implies x(\ell + 1) &= Sx(\ell). \end{aligned} \quad (\text{VII.15})$$

We get a similar expression for the discrete diffusion problem, which is usually called the *random walk* problem. Thus in this section, we will explore the following equations.

$$\begin{aligned} x(\ell + 1) &= Sx(\ell) \quad \text{with } x(0) = x_0 \quad \text{and} \\ p(\ell + 1) &= p(\ell)S \quad \text{with } p(0) = p_0. \end{aligned} \quad (\text{VII.16})$$

For the discrete processes, we have, as for the continuous ones, a convenient characterization of the asymptotic behavior. However, in the discrete case the solution $x(\ell) = S^\ell x_0$ might exhibit periodic behavior and so would not converge, as we shall see in some examples below. Thus we study the limit of the *average* instead

$$\frac{1}{\ell} \sum_{j=0}^{\ell-1} S^j x_0, \quad \text{as } \ell \rightarrow \infty. \quad (\text{VII.17})$$

If $S^\ell x_0$ does have a limit it will be same as the limit of the average. Using the average, we can proceed similarly to the continuous case in section VI.

Theorem VII.1 *Given a Laplacian $\mathcal{L} = (I - S)$ and Γ as in Definition V.3. We have*

$$\lim_{\ell \rightarrow \infty} \frac{1}{\ell} \sum_{j=0}^{\ell-1} S^j = \Gamma.$$

Proof: The proof is very similar to that of Theorem VI.1. The difference is in the analogue of equation (VI.13). This time, λ_i refers to an eigenvalue of S , not L , but with otherwise the same notation, instead of that equation, we now have

$$x(\ell) = S^\ell x(0) = \sum_{i=1}^n \gamma_i \lambda_i^\ell \xi_i(t). \quad (\text{VII.18})$$

For $i \in \{1, \dots, k\}$, we have $\lambda_i = 1$ and $\xi_i(t) = \alpha_i$. The left and right eigenspaces of the eigenvalue 1 are the same as the left and right kernels of the Laplacian (Theorem's V.1 and V.2).

By Gersgorin's theorem, the eigenvalues of S are in the closed unit ball. By Perron-Frobenius [8, 16], every eigenvalue λ not equal to 1 but with modulus 1 has equal algebraic and geometric multiplicity. Thus if v is the eigenvector corresponding to λ ,

$$\frac{1}{\ell} \sum_{j=0}^{\ell-1} \lambda^j v = 0.$$

All other eigenvalues have modulus less than 1, and so their contribution in the sum also vanishes. ■

Corollary VII.2 *The solutions $x(\ell)$ and $p(\ell)$ of (VII.16) satisfy:*

$$\lim_{\ell \rightarrow \infty} \frac{1}{\ell} \sum_{j=0}^{\ell-1} x(j) = \Gamma x_0 \quad \text{and} \quad \lim_{\ell \rightarrow \infty} \frac{1}{\ell} \sum_{j=0}^{\ell-1} p(j) = p_0 \Gamma.$$

The fact that in the discrete case, we have to account for periodic behavior explains why in the discrete case, we must take a limit of an *average*, while in the continuous case, it is sufficient to just take a limit (see Theorem VI.1). Again, taking Figure II.1 as example with \mathcal{L} given in equation (III.5), consider both discrete diffusion and discrete random walk with initial condition concentrated on vertex 7 only. As in Section VI, we get

$$\begin{aligned} \lim_{\ell \rightarrow \infty} \frac{1}{\ell} \sum_{j=0}^{\ell-1} x(j) &= \mathbf{0} \quad \text{and} \\ \lim_{\ell \rightarrow \infty} \frac{1}{\ell} \sum_{j=0}^{\ell-1} p(j) &= \frac{1}{9} (3, 0, 2, 2, 2, 0, 0). \end{aligned}$$

Notice that the random walker, when it arrives at vertex 3, undergoes periodic behavior. We effectively take the average of that behavior.

To isolate this periodic behavior, take the subgraph formed by the three vertices 3, 4 and 5 of the graph in figure II.1. This forms a cycle graph of order 3. The asymptotic behavior of the discrete solutions $x(\ell) = S^\ell x(0)$ are determined by the eigenvalues of S that have modulus equal to 1, because all other terms in (VII.18) tend to 0 (since the associated eigenvalues have modulus less than 1). These are exactly the eigenvalues of the submatrix of S in (VII.18) restricted to vertices 3, 4, and 5. Notice that the problem of periodic behavior “disappears” in the continuous system, because there we consider the eigenvalues of $-\mathcal{L} = S - I$. So all eigenvalues shift to the left, and all but one now have negative real part. This is illustrated in Figures VII.4 and VII.5.




FIG. VII.4. Eigenvalues of S




FIG. VII.5. Eigenvalues of $-\mathcal{L} = S - I$

Finally, we briefly discuss an alternative to the naive discretization we have been studying in this section so far. This is the so-called time one map of the continuous time dynamics of equation VI.11.

$$x(\ell) = e^{-\mathcal{L}} x(\ell - 1) = (S^d)^\ell x(0),$$

where we define $S^d = e^{-\mathcal{L}}$. We show that $S^{(d)}$ is a row stochastic matrix. We start with two expansions of that

matrix.

$$\begin{aligned} S^d &= e^{-\mathcal{L}} = \left(I - \mathcal{L} + \frac{\mathcal{L}^2}{2} - \dots \right), \\ S^d &= e^{S-I} = e^{-1} \left(I + S + \frac{S^2}{2} + \dots \right). \end{aligned} \quad (\text{VII.19})$$

From the first of these equations one easily derives that the matrix S^d has row sum one (just right multiply by the vector $\mathbf{1}$). The second equation implies that all its entries are non-negative. Therefore, $S^d = e^{-\mathcal{L}}$ is a row stochastic matrix.

It is clear that given any Laplacian \mathcal{L} we can (in theory) always compute its time one map $e^{-\mathcal{L}}$. It is interesting that the opposite is not true. From the second expansion in (VII.19), one can deduce that for every pair of vertices (i, j) in the graph associated to the weighted adjacency matrix S^d such that there is a path $i \rightsquigarrow j$, there is an edge $i \rightarrow j$, though its weight might be very small. (A graph with this property is called *transitively closed*.) Among other things, this implies that no time one map of a Laplacian can generate periodic behavior. In addition no Laplacian can generate a time one map with a zero eigenvalue.

VIII. CONCLUDING REMARKS

We analyzed the asymptotic behavior of general first order Laplacian processes on digraphs. The most important of these are diffusion and consensus with both continuous and discrete time. We have seen that diffusion and consensus are *dual* processes.

We remark here that given a continuous time diffusion or consensus process, it is always possible to find its time 1 map. But vice versa is not always possible. The reason is evident from the second part of (VII.19). That equation shows that any in time one map, every edge is realized, though not with the same weight. More details are given in [13].

The theory presented here has more applications than anyone can write down. We mentioned a few in the introduction. Here we want to mention briefly a few specific uses of the algorithms derived here. The first is that the duality described here can be used to give a new interpretation of the famed Pagerank algorithm as described in [8]. The interpretation is that the Pagerank of a site corresponds to the influence of the owner managing that site. For details see [13].

We end with two “folklore” results that can be easily proved with the tools of this paper. G is a (weakly connected) digraph with rw Laplacian \mathcal{L} . The union of its cabals is called B . Its complement is denoted as B^c . First, a random walker starting at vertex j has a probability $\gamma_{m,j}$ of ending up in the m th cabal B_m . The second result is that expected hitting time $\tau(i)$ for a random walk starting at vertex i to reach (or hit) B , is the unique solution of

$$\mathcal{L}\tau = \mathbf{1}_{B^c} \text{ with } \tau|_B = 0.$$

-
- [1] A. Broder, R. Kumar, F. Maghoul, P. Raghavan, S. Rajagopalan, R. Stata, A. Tomkins, and J. Wiener, Graph structure in the web, *Computer Networks* **33**, 309 (2000).
- [2] P. Carrington, J. Scott, and S. Wasserman, *Models and Methods in Social Network Analysis* (Cambridge University Press, 2005).
- [3] R. M. May, Qualitative stability in model ecosystems, *Ecology* **54**, 638 (1973).
- [4] T. Jombart, R. M. Eggo, P. J. Dodd, and F. Balloux, Reconstructing disease outbreaks from genetic data: a graph approach, *Heredity* **106**, 383 (2010).
- [5] S. Rao, A. van der Schaft, and B. Jayawardhana, A graph-theoretical approach for the analysis and model reduction of complex-balanced chemical reaction networks, *Journal of Mathematical Chemistry* **51**, 2401 (2013).
- [6] R. Angles and C. Gutierrez, Survey of graph database models, *ACM Computing Surveys* **40**, 1 (2008).
- [7] R. Ahlswede, N. Cai, S. R. Li, and R. W. Yeung, Network information flow, *IEEE Transactions on Information Theory* **46**, 1204 (2000).
- [8] S. Sternberg, *Dynamical systems* (Dover, Mineola, NY, 2010).
- [9] J. A. Fax and R. M. Murray, Information flow and cooperative control of vehicle formations, *IFAC Proceedings Volumes* **35**, 115 (2002).
- [10] C. Godsil and G. Royle, *Algebraic Graph Theory* (Springer, 2001).
- [11] F. R. K. Chung, *Spectral Graph Theory* (American Mathematical Society, 1997).
- [12] J. S. Caughman and J. J. P. Veerman, Kernels of directed graph laplacians, *The Electronic Journal of Combinatorics [electronic only]* **13**, Research paper R39 (2006).
- [13] J. J. P. Veerman and E. Kummel, Diffusion and consensus on weakly connected directed graphs, *Linear Algebra and its Applications* , 184 (2019).
- [14] J. J. P. Veerman, Digraphs ii: Diffusion and consensus on digraphs, <https://www.sci.unich.it/mmcs2019/slides/2019-Digraphs2.pdf> (2019).
- [15] R. A. Horn and C. R. Johnson, *Matrix analysis*, 2nd ed. (Cambridge University Press, New York, 2017).
- [16] M. Boyle, Notes on perron-frobenius theory of nonnegative matrices, www.math.umd.edu/~mboyle/courses/475sp05/spec.pdf, accessed 2019-10-26.
- [17] V. I. Arnold, *Ordinary Differential Equations*, 3rd ed. (Springer, Heidelberg, Berlin, 1992).